

COX8C Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55382**Specification**

COX8C Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	Q7Z4L0
Host	Rabbit
Clonality	Polyclonal
Calculated MW	8129

COX8C Polyclonal Antibody - Additional Information**Gene ID** 341947**Other Names**

Cytochrome c oxidase subunit 8C, mitochondrial, Cytochrome c oxidase polypeptide 8 isoform 3, Cytochrome c oxidase polypeptide VIII isoform 3, COX VIII-3, Cytochrome c oxidase subunit 8-3, COX8-3, Cytochrome c oxidase subunit VIIC, COX8C

Format

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

COX8C Polyclonal Antibody - Protein Information**Name** COX8C**Function**

Component of the cytochrome c oxidase, the last enzyme in the mitochondrial electron transport chain which drives oxidative phosphorylation. The respiratory chain contains 3 multisubunit complexes succinate dehydrogenase (complex II, CII),

ubiquinol- cytochrome c oxidoreductase (cytochrome b-c1 complex, complex III, CIII) and cytochrome c oxidase (complex IV, CIV), that cooperate to transfer electrons derived from NADH and succinate to molecular oxygen, creating an electrochemical gradient over the inner membrane that drives transmembrane transport and the ATP synthase.

Cytochrome c oxidase is the component of the respiratory chain that catalyzes the reduction of oxygen to water. Electrons originating from reduced cytochrome c in the intermembrane space (IMS) are transferred via the dinuclear copper A center (CU(A)) of subunit 2 and heme A of subunit 1 to the active site in subunit 1, a binuclear center (BNC) formed by heme A3 and copper B (CU(B)). The BNC reduces molecular oxygen to 2 water molecules using 4 electrons from cytochrome c in the IMS and 4 protons from the mitochondrial matrix.

Cellular Location

Mitochondrion inner membrane
{ECO:0000250|UniProtKB:P10175};
Single-pass membrane protein
{ECO:0000250|UniProtKB:P10175}

Tissue Location

It is not yet known where COX8C is expressed.

COX8C Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)