

DCTN5 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55516

Specification

DCTN5 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q9BTE1
Reactivity	Rat, Pig, Cow
Host	Rabbit
Clonality	Polyclonal
Calculated MW	20127

DCTN5 Polyclonal Antibody - Additional Information

Gene ID 84516

Other Names


Dynactin subunit 5, Dynactin subunit p25, DCTN5

Format

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glycerol

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.



Tissue/cell: rat kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-DCTN5 Polyclonal Antibody, Unconjugated(bs-14308R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

DCTN5 Polyclonal Antibody - Protein Information

Name DCTN5

Cellular Location

Cytoplasm, cytoskeleton. Chromosome, centromere, kinetochore

DCTN5 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)