

**DNAJB7 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP55543**

**Specification**

**DNAJB7 Polyclonal Antibody - Product Information**

Application	<b>IHC-P</b>
Primary Accession	<a href="#">Q7Z6W7</a>
Reactivity	<b>Rat, Dog, Cow</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>35434</b>

**DNAJB7 Polyclonal Antibody - Additional Information**

**Gene ID** 150353

**Other Names**

Dnaj homolog subfamily B member 7,  
DNAJB7, HSC3

**Format**

0.01M TBS(pH7.4) with 1% BSA, 0.09%  
(W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated  
freeze/thaw cycles. When reconstituted in  
sterile pH 7.4 0.01M PBS or diluent of  
antibody the antibody is stable for at least  
two weeks at 2-4 °C.

**DNAJB7 Polyclonal Antibody - Protein Information**

**Name** DNAJB7


**Synonyms** HSC3

**Function**

Probably acts as a co-chaperone.

**DNAJB7 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you  
may find useful for product applications.



Tissue/cell: mouse embryo tissue; 4%  
Paraformaldehyde-fixed and  
paraffin-embedded;

Antigen retrieval: citrate buffer ( 0.01M, pH  
6.0 ), Boiling bathing for 15min; Block  
endogenous peroxidase by 3% Hydrogen  
peroxide for 30min; Blocking buffer (normal  
goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-DNAJB7 Polyclonal  
Antibody, Unconjugated(bs-14374R) 1:200,  
overnight at 4°C, followed by conjugation to  
the secondary antibody(SP-0023) and  
DAB(C-0010) staining

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)