

**FCRL1 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56100****Specification**

---

**FCRL1 Polyclonal Antibody - Product Information**

Application	<b>WB, IHC-P, IHC-F, IF, ICC</b>
Primary Accession	<a href="#">Q96LA6</a>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>46936</b>

**FCRL1 Polyclonal Antibody - Additional Information****Gene ID** 115350**Other Names**

Fc receptor-like protein 1, FcR-like protein 1, FcRL1, Fc receptor homolog 1, FcRH1, IFGP family protein 1, hIFGP1, Immune receptor translocation-associated protein 5, CD307a, FCRL1, FCRH1, IFGP1, IRTA5

**Format**

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**FCRL1 Polyclonal Antibody - Protein Information****Name** FCRL1**Synonyms** FCRH1, IFGP1, IRTA5**Function**

May function as an activating coreceptor in B-cells. May function in B-cells activation and differentiation.

**Cellular Location**

Cell membrane; Single-pass type I

membrane protein

**Tissue Location**

Primarily expressed in secondary lymphoid tissues by mature subsets of B-cells. Detected in spleen, lymph node, heart, skeletal muscle, kidney, liver and placenta. Specifically expressed by mature B lineage cells with higher expression in naive versus memory B- cells (at protein level).

**FCRL1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)