

**GTF2IRD2 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56228**

**Specification**

**GTF2IRD2 Polyclonal Antibody - Product Information**

Application	<b>IHC-P, WB</b>
Primary Accession	<a href="#">Q86UP8</a>
Reactivity	<b>Rat, Pig, Dog, Cow</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>107168</b>

**GTF2IRD2 Polyclonal Antibody - Additional Information**

**Gene ID** 84163

**Other Names**

General transcription factor II-I repeat domain-containing protein 2A, GTF2I repeat domain-containing protein 2A, Transcription factor GTF2IRD2-alpha, GTF2IRD2, GTF2IRD2A

**Format**

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**GTF2IRD2 Polyclonal Antibody - Protein Information**

**Name** GTF2IRD2


**Synonyms** GTF2IRD2A

**Cellular Location**


Nucleus.

**Tissue Location**

Ubiquitous..



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GTF2IRD2) Polyclonal Antibody, Unconjugated (bs-16355R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample: K562 Cell (Human) Lysate at 40 ug  
 Primary: Anti-GTF2IRD2 (bs-16355R) at 1/300 dilution

## GTF2IRD2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 107 kD  
Observed band size: 107 kD