

**RASL11A Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP56836****Specification**

---

**RASL11A Polyclonal Antibody - Product Information**

Application	<b>IHC-P, IHC-F, IF, ICC</b>
Primary Accession	<a href="#">O6T310</a>
Reactivity	<b>Rat, Pig, Dog, Cow</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>27006</b>

**RASL11A Polyclonal Antibody - Additional Information****Gene ID** 387496**Other Names**

Ras-like protein family member 11A, 3.6.5.2, RASL11A ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=23802](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=23802))  
target="\_blank">HGNC:23802</a>)

**Format**

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**RASL11A Polyclonal Antibody - Protein Information****Name** RASL11A ([HGNC:23802](#))**Function**

Regulator of rDNA transcription. Acts in cooperation UBF/UBTF and positively regulates RNA polymerase I transcription (By similarity).

**Cellular Location**

Nucleus, nucleolus. Note=Associates with rDNA transcription unit throughout the cell cycle.

**Tissue Location**

Widely expressed. Down-regulated in prostate tumors compared to normal prostate tissue. High levels found in colon tumor and normal colon tissue followed by small intestine, liver, jejunum, ileum, bladder and aorta. Lowest levels observed in endothelial cells

**RASL11A Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)