

EBPL Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5735B

Specification

EBPL Antibody (C-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	O9BY08
Other Accession	NP_115954.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	23204
Antigen Region	177-206

EBPL Antibody (C-term) - Additional Information

Gene ID 84650

Other Names

Emopamil-binding protein-like,
Emopamil-binding-related protein, EBPL,
EBRP, ERP

Target/Specificity

This EBPL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 177-206 amino acids from the C-terminal region of human EBPL.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



EBPL Antibody (C-term) (Cat. #AP5735b) western blot analysis in NCI-H460 cell line lysates (15ug/lane). This demonstrates the EBPL antibody detected the EBPL protein (arrow).



EBPL Antibody (C-term) (Cat. #AP5735b) immunohistochemistry analysis in formalin fixed and paraffin embedded human lung carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the EBPL Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

EBPL Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

EBPL Antibody (C-term) - Protein Information

Name EBPL

Synonyms EBRP, ERP

Function

Does not possess sterol isomerase activity and does not bind sigma ligands.

Cellular Location

Endoplasmic reticulum membrane;
Multi-pass membrane protein


Tissue Location

Widely expressed with highest levels in liver, lung and kidney.

EBPL Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



EBPL Antibody (C-term) (Cat. #AP5735b) flow cytometric analysis of NCI-H460 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

EBPL Antibody (C-term) - Background

EBPL does not possess sterol isomerase activity and does not bind sigma ligands.

EBPL Antibody (C-term) - References

Ma, J., et al. Atherosclerosis
191(1):63-72(2007)
Moebius, F.F., et al. Biochem. J. 374 (PT 1),
229-237 (2003) :