

**ARHGAP20 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58143**

**Specification**

**ARHGAP20 Polyclonal Antibody - Product Information**

Application	<b>IHC-P</b>
Primary Accession	<a href="#">Q9P2F6</a>
Reactivity	<b>Rat, Dog, Cow</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>132608</b>

**ARHGAP20 Polyclonal Antibody - Additional Information**

**Gene ID** 57569

**Other Names**

Rho GTPase-activating protein 20, Rho-type GTPase-activating protein 20, ARHGAP20, KIAA1391

**Format**

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**ARHGAP20 Polyclonal Antibody - Protein Information**

**Name** ARHGAP20


**Synonyms** KIAA1391

**Function**

GTPase activator for the Rho-type GTPases by converting them to an inactive GDP-bound state.

**Tissue Location**

Expressed predominantly in the brain. Lower expression is found in lymph nodes.



Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ARHGAP20) Polyclonal Antibody, Unconjugated (bs-4271R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

## **ARHGAP20 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)