

**p53AIP1 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58591**

## Specification

### p53AIP1 Polyclonal Antibody - Product Information

Application	<b>IHC-P</b>
Primary Accession	<a href="#">Q9HCN2</a>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>12935</b>

### p53AIP1 Polyclonal Antibody - Additional Information

**Gene ID 63970**

#### Other Names

p53-regulated apoptosis-inducing protein 1,  
p53AIP1, TP53AIP1

#### Format

0.01M TBS(pH7.4) with 1% BSA, 0.09%  
(W/V) sodium azide and 50% Glyce

#### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

### p53AIP1 Polyclonal Antibody - Protein Information

**Name** TP53AIP1

#### Function


May play an important role in mediating p53/TP53-dependent apoptosis.

#### Cellular Location

Mitochondrion.

#### Tissue Location

Only found to be expressed in thymus.



Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-p53AIP1 Polyclonal Antibody, Unconjugated(bs-7094R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

## **p53AIP1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)