

SRRM4 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP58859

Specification

SRRM4 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F,
	IF
Primary Accession	<u>A7MD48</u>
Reactivity	Rat, Dog, Cow
Host	Rabbit
Clonality	Polyclonal
Calculated MW	68559

SRRM4 Polyclonal Antibody - Additional Information

Gene ID 84530

Other Names

Serine/arginine repetitive matrix protein 4, Medulloblastoma antigen MU-MB-2.76, Neural-specific serine/arginine repetitive splicing factor of 100 kDa, Neural-specific SR-related protein of 100 kDa, nSR100, SRRM4, KIAA1853

Format

0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SRRM4 Polyclonal Antibody - Protein Information

Name SRRM4

Synonyms KIAA1853

Function

Splicing factor specifically required for neural cell differentiation. Acts in conjunction with nPTB/PTBP2 by binding directly to its regulated target transcripts and promotes neural-specific exon inclusion in many genes that function in neural cell differentiation. Required to promote the inclusion of neural-specific exon 10 in nPTB/PTBP2, leading to increased expression of neural-specific nPTB/PTBP2. Also promotes the inclusion of exon 16 in DAAM1 in neuron extracts (By similarity). Promotes alternative splicing of REST transcripts to produce REST isoform 3 (REST4) with greatly reduced repressive activity, thereby activating expression of REST targets in neural cells (PubMed: <a hre f="http://www.uniprot.org/citations/306846 77" target=" blank">30684677). Plays an important role during embryonic development as well as in the proper functioning of the adult nervous system. Regulates alternative splicing events in genes with important neuronal functions (By similarity).

Cellular Location Nucleus {ECO:0000250|UniProtKB:Q8BKA3}.

Tissue Location Specifically expressed in neuronal cells (at protein level). Expressed in the cerebellum

SRRM4 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>