

C13orf18 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5893c

Specification

C13orf18 Antibody (Center) - Product Information

Application	WB, FC,E
Primary Accession	Q9H714
Other Accession	NP_079389.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	73457
Antigen Region	379-405

C13orf18 Antibody (Center) - Additional Information

Gene ID 80183

Other Names

Uncharacterized protein KIAA0226-like, KIAA0226L, C13orf18

Target/Specificity

This C13orf18 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 379-405 amino acids from the Central region of human C13orf18.

Dilution

WB~~1:1000
 FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

C13orf18 Antibody (Center) is for research



Anti-C13orf18 Antibody (Center) at 1:1000 dilution + Jurkat whole cell lysate
 Lysates/proteins at 20 µg per lane.
 Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.
 Predicted band size : 73 kDa
 Blocking/Dilution buffer: 5% NFDM/TBST.



C13orf18 Antibody (Center) (Cat. #AP5893c) flow cytometric analysis of Ramos cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

use only and not for use in diagnostic or therapeutic procedures.

C13orf18 Antibody (Center) - Protein Information

Name RUBCNL ([HGNC:20420](#))

Function

Regulator of autophagy that promotes autophagosome maturation by facilitating the biogenesis of phosphatidylinositol 3-phosphate (PtdIns(3)P) in late steps of autophagy (PubMed:[28306502](http://www.uniprot.org/citations/28306502)), PubMed:[30704899](http://www.uniprot.org/citations/30704899)). Acts by antagonizing RUBCN, thereby stimulating phosphatidylinositol 3-kinase activity of the PI3K/PI3KC3 complex (PubMed:[28306502](http://www.uniprot.org/citations/28306502)). Following anchorage to the autophagosomal SNARE STX17, promotes the recruitment of PI3K/PI3KC3 and HOPS complexes to the autophagosome to regulate the fusion specificity of autophagosomes with late endosomes/lysosomes (PubMed:[28306502](http://www.uniprot.org/citations/28306502)). Binds phosphoinositides phosphatidylinositol 3-phosphate (PtdIns(3)P), 4-phosphate (PtdIns(4)P) and 5-phosphate (PtdIns(5)P) (PubMed:[28306502](http://www.uniprot.org/citations/28306502)). In addition to its role in autophagy, acts as a regulator of lipid and glycogen homeostasis (By similarity). May act as a tumor suppressor (Probable).

Cellular Location

Cytoplasmic vesicle, autophagosome membrane; Peripheral membrane protein. Note=Associates with late autophagic structure (PubMed:28306502). Recruitment to autophagosome membrane is promoted by autophagic stimuli (PubMed:28306502)

Tissue Location

Expressed weakly in cervical carcinoma cell lines.

C13orf18 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)