

Wee1 (phospho Ser642) Polyclonal Antibody Catalog # AP67322

Specification

Wee1 (phospho Ser642) Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P30291
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Wee1 (phospho Ser642) Polyclonal Antibody - Additional Information

Gene ID 7465

Other Names

WEE1; Wee1-like protein kinase; WEE1hu;
Wee1A kinase

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA:
1/5000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5%
BSA and 0.02% sodium azide.

Storage Conditions


-20°C

Wee1 (phospho Ser642) Polyclonal Antibody - Protein Information

Name WEE1

Function

Acts as a negative regulator of entry into mitosis (G2 to M transition) by protecting the nucleus from cytoplasmically activated cyclin B1-complexed CDK1 before the onset of mitosis by mediating phosphorylation of CDK1 on 'Tyr-15'. Specifically phosphorylates and inactivates cyclin B1-complexed CDK1 reaching a maximum during G2 phase and a minimum as cells enter M phase. Phosphorylation of cyclin B1-CDK1 occurs exclusively on 'Tyr-15' and



Wee1 (phospho Ser642) Polyclonal Antibody - Background

Acts as a negative regulator of entry into mitosis (G2 to M transition) by protecting the nucleus from cytoplasmically activated cyclin B1-complexed CDK1 before the onset of mitosis by mediating phosphorylation of CDK1 on 'Tyr-15'. Specifically phosphorylates and inactivates cyclin B1-complexed CDK1 reaching a maximum during G2 phase and a minimum as cells enter M phase. Phosphorylation of cyclin B1-CDK1 occurs exclusively on 'Tyr-15' and phosphorylation of monomeric CDK1 does not occur. Its activity increases during S and G2 phases and decreases at M phase when it is hyperphosphorylated. A correlated decrease in protein level occurs at M/G1 phase, probably due to its degradation.

phosphorylation of monomeric CDK1 does not occur. Its activity increases during S and G2 phases and decreases at M phase when it is hyperphosphorylated. A correlated decrease in protein level occurs at M/G1 phase, probably due to its degradation.

Cellular Location

Nucleus.

Wee1 (phospho Ser642) Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)