

COLEC11 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP6840a

Specification

COLEC11 Antibody (N-term) - Product Information

Application WB, IHC-P, FC,E
Primary Accession
Reactivity WB, IHC-P, FC,E

O9BWP8
Human

Host Rabbit
Clonality Polyclonal
Isotype Rabbit Ig
Antigen Region 50-78

COLEC11 Antibody (N-term) - Additional Information

Gene ID 78989

Other Names

Collectin-11, Collectin kidney protein 1, CL-K1, COLEC11

Target/Specificity

This COLEC11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 50-78 amino acids from the N-terminal region of human COLEC11.

Dilution

WB~~1:1000 IHC-P~~1:50~100 FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

COLEC11 Antibody (N-term) is for research use only and not for use in diagnostic or



Western blot analysis of COLEC11 Antibody (N-term) (Cat. #AP6840a) in MDA-MB231 cell line lysates (35ug/lane). COLEC11 (arrow) was detected using the purified Pab.



Western blot analysis of COLEC11 (arrow) using rabbit polyclonal COLEC11 Antibody (N-term) (Cat. #AP6840a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the COLEC11 gene.



therapeutic procedures.

COLEC11 Antibody (N-term) - Protein Information

Name COLEC11

Function

Lectin that plays a role in innate immunity, apoptosis and embryogenesis (PubMed:23954398, PubMed:<a href="http://www.uniprot.org/citations/25912189"

target=" blank">25912189,

PubMed:<a href="http://www.uniprot.org/ci tations/21258343"

target="_blank">21258343).

Calcium-dependent lectin that binds self and non-self glycoproteins presenting high mannose oligosaccharides with at least one terminal alpha-1,2-linked mannose epitope (PubMed:<a href="http://www.uniprot.org/c itations/25912189"

target="_blank">25912189). Primarily recognizes the terminal disaccharide of the glycan (PubMed:<a href="http://www.uniprot.org/citations/25912189"

target="_blank">25912189). Also recognizes a subset of fucosylated glycans and lipopolysaccharides (PubMed:17179669,


PubMed:<a href="http://www.uniprot.org/ci tations/25912189"

target="_blank">25912189). Plays a role in innate immunity through its ability to bind non-self sugars presented by microorganisms and to activate the complement through the recruitment of MAPS1 (PubMed:<a href="http://www.uniprot.org/citations/20956340"


target="_blank">20956340,

PubMed: <a href="http://www.uniprot.org/ci tations/25912189"

target="_blank">25912189). Also plays a role in apoptosis through its ability to bind in a calcium-independent manner the DNA present at the surface of apoptotic cells and to activate the complement in response to this binding (Probable). Finally, plays a role in development, probably serving as a guidance cue during the migration of neural crest cells and other cell types during embryogenesis (PubMed:<a hr ef="http://www.uniprot.org/citations/21258343" target="_blank">21258343,



Formalin-fixed and paraffin-embedded human hepatocarcinoma with COLEC11 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.




Flow cytometric analysis of MDA-231 cells using COLEC11 Antibody (N-term)(bottom histogram) compared to a negative control (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

COLEC11 Antibody (N-term) - Background

COLEC11 is a member of the collectin family of C-type lectins, which contain a collagen-like domain and a carbohydrate recognition domain, and play a role in host-defense

COLEC11 Antibody (N-term) - References

Oguri, M., et.al., Am. J. Hypertens. (2009)





PubMed:28301481).

Cellular Location Secreted.

Tissue Location

Ubiquitous (PubMed:17179669). Detected in adrenal gland, kidney, liver, ovaries and testis (at protein level) (PubMed:20956340).

COLEC11 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture