

IGHA1 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7456b

Specification

IGHA1 Antibody (C-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P01876
Other Accession	P01877
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	37655
Antigen Region	257-286

IGHA1 Antibody (C-term) - Additional Information

Other Names

Ig alpha-1 chain C region, IGHAI

Target/Specificity

This IGHAI antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 257-286 amino acids from the C-terminal region of human IGHAI.

Dilution

WB~~1:1000
 IHC-P~~1:50~100
 FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IGHAI Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.



Western blot analysis of IGHAI antibody (C-term) (Cat.#AP7456b) in CEM cell line lysates (35ug/lane). IGHAI (arrow) was detected using the purified Pab .



Anti-IgA Antibody (C-term) at 1:1000 dilution + A2058 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 38 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

IGHA1 Antibody (C-term) - Protein Information

Name IGHAI

{ECO:0000303|PubMed:11340299,
ECO:0000303|Ref.12}


Function

Constant region of immunoglobulin heavy chains. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulin-secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens (PubMed:22158414, PubMed:20176268). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed:17576170, PubMed:20176268). Ig alpha is the major immunoglobulin class in body secretions (PubMed:2241915).


Cellular Location

Secreted. Cell membrane

IGHA1 Antibody (C-term) - Protocols



Formalin-fixed and paraffin-embedded human Lymph reacted with IGHAI Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of CEM cells using IGHAI Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

IGHA1 Antibody (C-term) - Background

Ig alpha is the major immunoglobulin class in body secretions. The protein may serve both to defend against local infection and to prevent access of foreign antigens to the general immunologic system.

IGHA1 Antibody (C-term) - References

Flanagan J.G., Lefranc M.-P. Cell
36:681-688(1984)
Putnam F.W., Liu Y.-S.V.J. Biol. Chem.
254:2865-2874(1979)

Provided below are standard protocols that you may find useful for product applications.

Hatzivassiliou G., Miller I. Immunity
14:277-289(2001)

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)