

**BUB1 (BUB1a) Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8058b**

**Specification**

**BUB1 (BUB1a) Antibody (C-term) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">O43683</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	1043-1073

**BUB1 (BUB1a) Antibody (C-term) - Additional Information**

Gene ID 699

**Other Names**

Mitotic checkpoint serine/threonine-protein kinase BUB1, hBUB1, BUB1A, BUB1, BUB1L

**Target/Specificity**

This BUB1 (BUB1a) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1043-1073 amino acids from the C-terminal region of human BUB1 (BUB1a).

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100

**Format**


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

BUB1 (BUB1a) Antibody (C-term) is for research use only and not for use in



BUB1A Antibody (F1058) (Cat.#AP8058b) western blot analysis in T47D cell line lysates (35ug/lane). This demonstrates the BUB1A antibody detected the BUB1A protein (arrow).



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

diagnostic or therapeutic procedures.

#### **BUB1 (BUB1a) Antibody (C-term) - Protein Information**

**Name** BUB1

**Synonyms** BUB1L

#### **Function**

Serine/threonine-protein kinase that performs 2 crucial functions during mitosis: it is essential for spindle-assembly checkpoint signaling and for correct chromosome alignment. Has a key role in the assembly of checkpoint proteins at the kinetochore, being required for the subsequent localization of CENPF, BUB1B, CENPE and MAD2L1. Required for the kinetochore localization of PLK1. Required for centromeric enrichment of AUKRB in prometaphase. Plays an important role in defining SGO1 localization and thereby affects sister chromatid cohesion. Acts as a substrate for anaphase-promoting complex or cyclosome (APC/C) in complex with its activator CDH1 (APC/C-Cdh1). Necessary for ensuring proper chromosome segregation and binding to BUB3 is essential for this function. Can regulate chromosome segregation in a kinetochore-independent manner. Can phosphorylate BUB3. The BUB1-BUB3 complex plays a role in the inhibition of APC/C when spindle-assembly checkpoint is activated and inhibits the ubiquitin ligase activity of APC/C by phosphorylating its activator CDC20. This complex can also phosphorylate MAD1L1. Kinase activity is essential for inhibition of APC/CCDC20 and for chromosome alignment but does not play a major role in the spindle-assembly checkpoint activity. Mediates cell death in response to chromosome missegregation and acts to suppress spontaneous tumorigenesis.

#### **Cellular Location**

Nucleus. Chromosome, centromere, kinetochore. Note=Nuclear in interphase cells. Accumulates gradually during G1 and S phase of the cell cycle, peaks at G2/M, and drops dramatically after mitosis. Localizes to the outer kinetochore. Kinetochore localization is required for normal mitotic timing and checkpoint response to spindle damage and occurs

#### **BUB1 (BUB1a) Antibody (C-term) - Background**

This gene encodes a kinase involved in spindle checkpoint function. The kinase functions in part by phosphorylating a member of the mitotic checkpoint complex and activating the spindle checkpoint. Mutations in this gene have been associated with aneuploidy and several forms of cancer.

#### **BUB1 (BUB1a) Antibody (C-term) - References**

- Shichiri, M., et al., *Cancer Res.* 62(1):13-17 (2002).  
Cayrol, C., et al., *Biochem. Biophys. Res. Commun.* 298(5):720-730 (2002).  
Nakagawa, H., et al., *Oncol. Rep.* 9(6):1229-1232 (2002).  
Ru, H.Y., et al., *Oncogene* 21(30):4673-4679 (2002).  
Lin, S.F., et al., *Leuk. Lymphoma* 43(2):385-391 (2002).

very early in prophase. AURKB, KNL1 and INCENP are required for kinetochore localization (By similarity)

**Tissue Location**

High expression in testis and thymus, less in colon, spleen, lung and small intestine. Expressed in fetal thymus, bone marrow, heart, liver, spleen and thymus. Expression is associated with cells/tissues with a high mitotic index

**BUB1 (BUB1a) Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)