

CECR5 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP8560c

Specification

CECR5 Antibody (Center) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	<u>Q9BXW7</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	46321
Antigen Region	164-190

CECR5 Antibody (Center) - Additional Information

Gene ID 27440

Other Names

Cat eye syndrome critical region protein 5, CECR5

Target/Specificity

This CECR5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 164-190 amino acids from the Central region of human CECR5.

Dilution

WB~~1:1000 IHC-P~~1:50~100 FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CECR5 Antibody (Center) is for research use only and not for use in diagnostic or



Western blot analysis of CECR5 Antibody (Center) (Cat. #AP8560c) in Jurkat cell line lysates (35ug/lane). CECR5 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human hepatocarcinoma reacted with CECR5 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



therapeutic procedures.

CECR5 Antibody (Center) - Protein Information


Name HDHD5 (HGNC:1843)

Tissue Location Widely expressed.

CECR5 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>



Flow cytometric analysis of jurkat cells using CECR5 Antibody (Center)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

CECR5 Antibody (Center) - References

Footz,T.K., et.al., Genome Res. 11 (6), 1053-1070 (2001)