

C5 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8577a**Specification****C5 Antibody (N-term) - Product Information**

Application	WB, IHC-P, FC,E
Primary Accession	P01031
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	253-281

C5 Antibody (N-term) - Additional Information**Gene ID 727****Other Names**

Complement C5, C3 and PZP-like alpha-2-macroglobulin domain-containing protein 4, Complement C5 beta chain, Complement C5 alpha chain, C5a anaphylatoxin, Complement C5 alpha' chain, C5, CPAMD4

Target/Specificity

This C5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 253-281 amino acids of human C5.

Dilution


WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Western blot analysis of C5 Antibody (N-term) (Cat. #AP8577a) in HL-60 cell line lysates (35ug/lane). C5 (arrow) was detected using the purified Pab.(2ug/ml)



Formalin-fixed and paraffin-embedded human breast carcinoma with C5 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

C5 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

C5 Antibody (N-term) - Protein Information

Name C5

Synonyms CPAMD4

Function

Activation of C5 by a C5 convertase initiates the spontaneous assembly of the late complement components, C5-C9, into the membrane attack complex. C5b has a transient binding site for C6. The C5b-C6 complex is the foundation upon which the lytic complex is assembled.

Cellular Location

Secreted.


C5 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C5 Antibody (N-term) - Citations

- [SAK-HV Triggered a Short-period Lipid-lowering Biotherapy Based on the Energy Model of Liver Proliferation via a Novel Pathway.](#)



Flow cytometric analysis of MDA-231 cells using C5 Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

C5 Antibody (N-term) - Background

C5 is the fifth component of complement, which plays an important role in inflammatory and cell killing processes. This protein is comprised of alpha and beta polypeptide chains that are linked by a disulfide bridge. An activation peptide, C5a, which is an anaphylatoxin that possesses potent spasmogenic and chemotactic activity, is derived from the alpha polypeptide via cleavage with a convertase. The C5b macromolecular cleavage product can form a complex with the C6 complement component, and this complex is the basis for formation of the membrane attack complex, which includes additional complement components.

C5 Antibody (N-term) - References

McClure,A., et.al., Rheumatology (Oxford) 48 (11), 1369-1374 (2009)
DiScipio,R.G.et.al., J. Biol. Chem. 267 (24), 17087-17094 (1992)