

CAT Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8623c

Specification

CAT Antibody (Center) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P04040
Other Accession	P04762 , O62839 , P24270 , Q9PT92 , P00432
Reactivity Predicted	Human, Mouse Bovine, Zebrafish, Pig, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	152-180

CAT Antibody (Center) - Additional Information

Gene ID 847

Other Names
Catalase, CAT

Target/Specificity

This CAT antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 152-180 amino acids from the Central region of human CAT.

Dilution

WB~~1:2000
IHC-P~~1:50~100
FC~~1:10~50


Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions



All lanes : Anti-CAT Antibody (Center) at 1:2000 dilution Lane 1: HepG2 whole cell lysate Lane 2: Mouse kidney lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 60 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human lung carcinoma reacted with CAT Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

CAT Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

CAT Antibody (Center) - Protein Information

Name CAT

Function

Occurs in almost all aerobically respiring organisms and serves to protect cells from the toxic effects of hydrogen peroxide. Promotes growth of cells including T-cells, B-cells, myeloid leukemia cells, melanoma cells, mastocytoma cells and normal and transformed fibroblast cells.

Cellular Location

Peroxisome.


CAT Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CAT Antibody (Center) - Citations

- [Cardiac fatty acid uptake and metabolism in the rat model of polycystic ovary syndrome.](#)
- [The expression and activity of antioxidant enzymes in the liver of rats exposed to high-fructose diet in period from weaning to adulthood.](#)



CAT Antibody (Center) - Background

CAT occurs in almost all aerobically respiring organisms and serves to protect cells from the toxic effects of hydrogen peroxide. It promotes growth of cells including T-cells, B-cells, myeloid leukemia cells, melanoma cells, mastocytoma cells and normal and transformed fibroblast cells.

CAT Antibody (Center) - References

Oh, J.H., et al., Mamm. Genome 16 (12), 942-954 (2005)