

LTF Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8698A

Specification

LTF Antibody - Product Information

Application	WB, FC,E
Primary Accession	P02788
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig

LTF Antibody - Additional Information

Gene ID 4057

Other Names

Lactotransferrin, Lactoferrin, 3421-,
Growth-inhibiting protein 12,
Talaktoferrin, Lactoferricin-H, Lfcin-H,
Kaliocin-1, Lactoferroxin-A, Lactoferroxin-B,
Lactoferroxin-C, LTF, GIG12, LF

Target/Specificity

This LTF antibody is generated from rabbits immunized with LTF recombinant protein.

Dilution

WB~~1:1000
FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

LTF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Western blot analysis of LTF Antibody (Cat. #AP8698a) in MDA-MB231 cell line lysates (35ug/lane). LTF (arrow) was detected using the purified Pab.



Western blot analysis of LTF Antibody (Cat. #AP8698a) in mouse spleen tissue lysates (35ug/lane). LTF (arrow) was detected using the purified Pab.

LTF Antibody - Protein Information

Name LTF

Synonyms GIG12, LF

Function

Transferrins are iron binding transport proteins which can bind two Fe(3+) ions in association with the binding of an anion, usually bicarbonate.

Cellular Location

[Isoform 1]: Secreted. Cytoplasmic granule. Note=Secreted into most exocrine fluids by various endothelial cells Stored in the secondary granules of neutrophils


Tissue Location

High levels are found in saliva and tears, intermediate levels in serum and plasma, and low levels in urine. In kidney, detected in the distal collecting tubules in the medulla but not in the cortical region or in blood vessels. Detected in peripheral blood neutrophils (at protein level). Isoform 1 and isoform DeltaLf are expressed in breast, prostate, spleen, pancreas, kidney, small intestine, lung, skeletal muscle, uterus, thymus and fetal liver Isoform 1 is expressed in brain, testis and peripheral blood leukocytes; isoform DeltaLf is barely detectable in these tissues Isoform DeltaLf is expressed in placenta, liver and ovary; isoform 1 is barely detectable in these tissues. In kidney, isoform 1 is expressed at high levels in the collecting tubules of the medulla but at very low levels in the cortex.


LTF Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)




All lanes : Anti-LTF Antibody at 1:4000 dilution Lane 1: A549 whole cell lysate Lane 2: Human plasma tissue lysate Lane 3: Human fetal kidney tissue lysate Lane 4: Human breast tissue lysate Lane 5: Human uterus tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 78 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



All lanes : Anti-LTF Antibody at 1:1000-2000 dilution Lane 1: Human plasma tissue lysate Lane 2: HeLa whole cell lysate Lane 3: Human uterus tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 78 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



All lanes : Anti-LTF Antibody at 1:1000 dilution
 Lane 1: Human plasma tissue lysate
 Lane 2: Human fetal kidney tissue lysate
 Lane 3: Human uterus tissue lysate
 Lane 4: Human skeletal muscle tissue lysate
 Lysates/proteins at 20 µg per lane.
 Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.
 Predicted band size : 78 kDa
 Blocking/Dilution buffer: 5% NFDM/TBST.



LTF Antibody (Cat. #AP8698a) flow cytometric analysis of MDA-MB231 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

LTF Antibody - Background

LTF is found in the secondary granules of

neutrophils. The protein is a major iron-binding protein in milk and body secretions with an antimicrobial activity, making it an important component of the non-specific immune system. The protein demonstrates a broad spectrum of properties, including regulation of iron homeostasis, host defense against a broad range of microbial infections, anti-inflammatory activity, regulation of cellular growth and differentiation and protection against cancer development and metastasis.

LTF Antibody - References

Willnow, T.E., et.al., J. Biol. Chem. 267 (36), 26172-26180 (1992)