

CDC42EP3 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8775a

Specification

CDC42EP3 Antibody (N-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q9UKI2
Other Accession	Q9CQC5 , NP_006440
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	1-30

CDC42EP3 Antibody (N-term) - Additional Information

Gene ID 10602

Other Names

Cdc42 effector protein 3, Binder of Rho GTPases 2, MSE55-related Cdc42-binding protein, CDC42EP3, BORG2, CEP3

Target/Specificity

This CDC42EP3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human CDC42EP3.

Dilution


WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw



Western blot analysis of CDC42EP3 Antibody (N-term) (Cat. #AP8775a) in 293 cell line lysates (35ug/lane). CDC42EP3 (arrow) was detected using the purified Pab.



Western blot analysis of CDC42EP3 (arrow) using rabbit polyclonal CDC42EP3 Antibody (N-term) (Cat. #AP8775a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the CDC42EP3 gene.

cycles.

Precautions

CDC42EP3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CDC42EP3 Antibody (N-term) - Protein Information

Name CDC42EP3

Synonyms BORG2, CEP3

Function

Probably involved in the organization of the actin cytoskeleton. May act downstream of CDC42 to induce actin filament assembly leading to cell shape changes. Induces pseudopodia formation in fibroblasts.

Cellular Location

Endomembrane system; Peripheral membrane protein. Cytoplasm, cytoskeleton


Tissue Location

Highly expressed in the heart and weakly in the brain.


CDC42EP3 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Formalin-fixed and paraffin-embedded human testis tissue reacted with CDC42EP3 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



CDC42EP3 Antibody (N-term) (Cat. #AP8775a) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

CDC42EP3 Antibody (N-term) - Background

CDC42EP3 is probably involved in the organization of the actin cytoskeleton. May act downstream of CDC42 to induce actin filament assembly leading to cell shape changes. It induces pseudopodia formation in fibroblasts.

CDC42EP3 Antibody (N-term) - References

Simpson, J.C., et al., EMBO Rep. 1 (3), 287-292 (2000)