

RNF8 Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9820a

Specification

RNF8 Antibody - Product Information

| | |
|-------------------|------------------------|
| Application | WB, FC,E |
| Primary Accession | 076064 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit Ig |
| Calculated MW | 55518 |

RNF8 Antibody - Additional Information

Gene ID 9025

Other Names

E3 ubiquitin-protein ligase RNF8, hRNF8, 632-, RING finger protein 8, RNF8, KIAA0646

Target/Specificity

This RNF8 antibody is generated from rabbits immunized with recombinant protein of human RNF8.

Dilution

WB~~1:1000
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.


Precautions

RNF8 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

RNF8 Antibody - Protein Information



Western blot analysis of RNF8 Antibody (Cat. #AP9820a) in Jurkat cell line lysates (35ug/lane). RNF8 (arrow) was detected using the purified Pab.



RNF8 Antibody (Cat. #AP9820a) flow cytometric analysis of Jurkat cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

RNF8 Antibody - Background

RNF8 contains a RING finger motif and a FHA

Name RNF8
{ECO:0000255|HAMAP-Rule:MF_03067}

Synonyms KIAA0646

Function

E3 ubiquitin-protein ligase that plays a key role in DNA damage signaling via 2 distinct roles: by mediating the 'Lys-63'-linked ubiquitination of histones H2A and H2AX and promoting the recruitment of DNA repair proteins at double-strand breaks (DSBs) sites, and by catalyzing 'Lys-48'-linked ubiquitination to remove target proteins from DNA damage sites. Following DNA DSBs, it is recruited to the sites of damage by ATM-phosphorylated MDC1 and catalyzes the 'Lys-63'-linked ubiquitination of histones H2A and H2AX, thereby promoting the formation of TP53BP1 and BRCA1 ionizing radiation-induced foci (IRIF). Also controls the recruitment of UIMC1-BRCC3 (RAP80-BRCC36) and PAXIP1/PTIP to DNA damage sites. Also recruited at DNA interstrand cross-links (ICLs) sites and catalyzes 'Lys-63'-linked ubiquitination of histones H2A and H2AX, leading to recruitment of FAAP20/C1orf86 and Fanconi anemia (FA) complex, followed by interstrand cross-link repair. H2A ubiquitination also mediates the ATM-dependent transcriptional silencing at regions flanking DSBs in cis, a mechanism to avoid collision between transcription and repair intermediates. Promotes the formation of 'Lys-63'-linked polyubiquitin chains via interactions with the specific ubiquitin-conjugating UBE2N/UBC13 and ubiquitinates non-histone substrates such as PCNA. Substrates that are polyubiquitinated at 'Lys-63' are usually not targeted for degradation. Also catalyzes the formation of 'Lys-48'-linked polyubiquitin chains via interaction with the ubiquitin-conjugating UBE2L6/UBCH8, leading to degradation of substrate proteins such as CHEK2, JMJD2A/KDM4A and KU80/XRCC5: it is still unclear how the preference toward 'Lys-48'- versus 'Lys-63'-linked ubiquitination is regulated but it could be due to RNF8 ability to interact with specific E2 specific ligases. For instance, interaction with phosphorylated HERC2 promotes the association between RNF8 and UBE2N/UBC13 and favors the specific

domain. This protein has been shown to interact with several class II ubiquitin-conjugating enzymes (E2), including UBE2E1/UBCH6, UBE2E2, and UBE2E3, and may act as an ubiquitin ligase (E3) in the ubiquitination of certain nuclear proteins.

RNF8 Antibody - References

- Noon, A.T., et al. *Nat. Cell Biol.* 12(2):177-184(2010)
Ramachandran, S., et al. *Proc. Natl. Acad. Sci. U.S.A.* 107(2):809-814(2010)
Marteijn, J.A., et al. *J. Cell Biol.* 186(6):835-847(2009)
Gong, Z., et al. *J. Biol. Chem.* 284(11):7284-7293(2009)
Zhang, S., et al. *Cell Cycle* 7(21):3399-3404(2008)

formation of 'Lys-63'- linked ubiquitin chains. Promotes non-homologous end joining (NHEJ) by promoting the 'Lys-48'-linked ubiquitination and degradation of KU80/XRCC5. Following DNA damage, mediates the ubiquitination and degradation of JMJD2A/KDM4A in collaboration with RNF168, leading to unmask H4K20me2 mark and promote the recruitment of TP53BP1 at DNA damage sites (PubMed:11322894, PubMed:14981089, PubMed:17724460, PubMed:18001824, PubMed:18001825, PubMed:18006705, PubMed:18077395, PubMed:18337245, PubMed:18948756, PubMed:19015238, PubMed:19124460, PubMed:19202061, PubMed:19203578, PubMed:19203579, PubMed:20550933, PubMed:21558560

target="_blank">>21558560,
PubMed:<a href="http://www.uniprot.org/citations/21857671"
target="_blank">>21857671,
PubMed:<a href="http://www.uniprot.org/citations/21911360"
target="_blank">>21911360,
PubMed:<a href="http://www.uniprot.org/citations/22266820"
target="_blank">>22266820,
PubMed:<a href="http://www.uniprot.org/citations/22373579"
target="_blank">>22373579,
PubMed:<a href="http://www.uniprot.org/citations/22531782"
target="_blank">>22531782,
PubMed:<a href="http://www.uniprot.org/citations/22705371"
target="_blank">>22705371,
PubMed:<a href="http://www.uniprot.org/citations/22865450"
target="_blank">>22865450,
PubMed:<a href="http://www.uniprot.org/citations/22980979"
target="_blank">>22980979).
Following DNA damage, mediates the ubiquitination and degradation of POLD4/p12, a subunit of DNA polymerase delta. In the absence of POLD4, DNA polymerase delta complex exhibits higher proofreading activity (PubMed:<a href="http://www.uniprot.org/citations/23233665"
target="_blank">>23233665). In addition to its function in damage signaling, also plays a role in higher-order chromatin structure by mediating extensive chromatin decondensation. Involved in the activation of ATM by promoting histone H2B ubiquitination, which indirectly triggers histone H4 'Lys-16' acetylation (H4K16ac), establishing a chromatin environment that promotes efficient activation of ATM kinase. Required in the testis, where it plays a role in the replacement of histones during spermatogenesis. At uncapped telomeres, promotes the joining of deprotected chromosome ends by inducing H2A ubiquitination and TP53BP1 recruitment, suggesting that it may enhance cancer development by aggravating telomere-induced genome instability in case of telomeric crisis. Promotes the assembly of RAD51 at DNA DSBs in the absence of BRCA1 and TP53BP1. Also involved in class switch recombination in immune system, via its role in regulation of DSBs repair. May be required for proper exit

from mitosis after spindle checkpoint activation and may regulate cytokinesis. May play a role in the regulation of RXRA-mediated transcriptional activity. Not involved in RXRA ubiquitination by UBE2E2 (PubMed:11322894, PubMed:14981089, PubMed:17724460, PubMed:18001824, PubMed:18001825, PubMed:18006705, PubMed:18077395, PubMed:18337245, PubMed:18948756, PubMed:19015238, PubMed:19124460, PubMed:19202061, PubMed:19203578, PubMed:19203579, PubMed:20550933, PubMed:21558560, PubMed:21857671, PubMed:<a href="http://www.uniprot.org/ci

tations/21911360"
target="_blank">21911360,
PubMed:<a href="http://www.uniprot.org/ci
tations/22266820"
target="_blank">22266820,
PubMed:<a href="http://www.uniprot.org/ci
tations/22373579"
target="_blank">22373579,
PubMed:<a href="http://www.uniprot.org/ci
tations/22531782"
target="_blank">22531782,
PubMed:<a href="http://www.uniprot.org/ci
tations/22705371"
target="_blank">22705371,
PubMed:<a href="http://www.uniprot.org/ci
tations/22865450"
target="_blank">22865450,
PubMed:<a href="http://www.uniprot.org/ci
tations/22980979"
target="_blank">22980979).

Cellular Location

Nucleus

{ECO:0000255|HAMAP-Rule:MF_03067,
ECO:0000269|PubMed:11322894,
ECO:0000269|PubMed:14981089,
ECO:0000269|PubMed:16215985,
ECO:0000269|PubMed:23233665}.

Cytoplasm

{ECO:0000255|HAMAP-Rule:MF_03067}.
Midbody {ECO:0000255|HAMAP-
Rule:MF_03067}. Chromosome, telomere
{ECO:0000255|HAMAP-Rule:MF_03067}
Note=Recruited at uncapped telomeres (By
similarity). Following DNA damage, such as
double-strand breaks, recruited to the sites
of damage (PubMed:18001824,
PubMed:18077395, PubMed:22266820,
PubMed:23233665) During prophase,
concomitant with nuclear envelope
breakdown, localizes throughout the cell,
with a dotted pattern. In telophase, again in
the nucleus and also with a discrete dotted
pattern in the cytoplasm. In late telophase
and during cytokinesis, localizes in the
midbody of the tubulin bridge joining the
daughter cells. Does not seem to be
associated with condensed chromosomes at
any time during the cell cycle. During
spermatogenesis, sequestered in the
cytoplasm by PIWIL1: RNF8 is released
following ubiquitination and degradation of
PIWIL1
{ECO:0000255|HAMAP-Rule:MF_03067,
ECO:0000269|PubMed:18001824,
ECO:0000269|PubMed:18077395,
ECO:0000269|PubMed:22266820,

ECO:0000269|PubMed:23233665}

Tissue Location

Ubiquitous. In fetal tissues, highest expression in brain, thymus and liver. In adult tissues, highest levels in brain and testis, lowest levels in peripheral blood cells

RNF8 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)