

ASAH2 Antibody
Catalog # **ASC10741**

Specification

ASAH2 Antibody - Product Information

Application	WB
Primary Accession	O9NR71
Other Accession	NP_063946 , 221218981
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	ASAH2 antibody can be used for detection of ASAH2 by Western blot at 1 and 2 µg/mL.

ASAH2 Antibody - Additional Information

Gene ID	56624
Target/Specificity	ASAH2;

Reconstitution & Storage

ASAH2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

ASAH2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


ASAH2 Antibody - Protein Information

Name ASAH2

Synonyms HNAC1

Function

Plasma membrane ceramidase that hydrolyzes sphingolipid ceramides into



ASAH2 Antibody - Background

ASAH2 Antibody: Sphingolipids are hydrolyzed by ceramidases to yield sphingosine and fatty acids. These ceramidases are classified according to the pH range that supports their optimal activity. ASAH2 is a neutral ceramidase and key regulator of sphingolipid signaling metabolites at the cell surface, catalyzing the hydrolysis of the N-acyl linkage of ceramide at an optimal pH of 6.5-8.5. ASAH2 is a type II integral membrane protein that can be cleaved to yield a soluble secreted protein and acts as a repressor of apoptosis both by reducing C16-ceramide, thereby preventing ceramide-induced apoptosis, and generating sphingosine. Sphingosine exerts both mitogenic and apoptosis-inducing activities, and its phosphorylated form functions as an intra- and intercellular second messenger. ASAH2 is ubiquitously expressed primarily expressed with higher levels in the intestine, kidney, skeletal muscle and heart. Recent studies indicate that ASAH2 encoded neutral ceramidase is a key enzyme for the catabolism

sphingosine and free fatty acids at neutral pH (PubMed:10781606, PubMed:16229686, PubMed:26190575). Ceramides, sphingosine, and its phosphorylated form sphingosine-1-phosphate are bioactive lipids that mediate cellular signaling pathways regulating several biological processes including cell proliferation, apoptosis and differentiation (PubMed:15946935, PubMed:19345744, PubMed:24798654). Also catalyzes the reverse reaction allowing the synthesis of ceramides from fatty acids and sphingosine (PubMed:11278489, PubMed:17475390). Together with sphingomyelinase, participates in the production of sphingosine and sphingosine-1-phosphate from the degradation of sphingomyelin, a sphingolipid enriched in the plasma membrane of cells (PubMed:16061940). Also participates in the hydrolysis of ceramides from the extracellular milieu allowing the production of sphingosine-1-phosphate inside and outside cells (By similarity). This is the case for instance with the digestion of dietary sphingolipids in the intestinal tract (By similarity).

Cellular Location

[Neutral ceramidase]: Cell membrane;
Single-pass type II membrane protein
{ECO:0000250|UniProtKB:Q91XT9}.
Membrane raft
{ECO:0000250|UniProtKB:Q9JHE3};
Single-pass type II membrane protein
{ECO:0000250|UniProtKB:Q91XT9}.

of dietary sphingolipids and regulates the levels of bioactive sphingolipid metabolites in the intestinal tract.

ASAH2 Antibody - References

Nilsson A and Duan RD. Alkaline sphingomyelinases and ceramidases of the gastrointestinal tract. *Chem. Phys. Lipids*1999; 102:97-105.
Tani M, Okino N, Mitsutake S, et al. Molecular Cloning of the Full-length cDNA Encoding Mouse Neutral Ceramidase. *J. Biol. Chem.*2000; 275:11229-34.
Tani M, Iida H, and Ito M. O-Glycosylation of Mucin-like Domain Retains the Neutral Ceramidase on the Plasma Membranes as a Type II Integral Membrane Protein. *J. Biol. Chem.*2003; 278:10523-30.
Osawa Y, Uchinami H, Bielawski J, et al. Roles for C16-ceramide and sphingosine 1-phosphate in regulating hepatocyte apoptosis in response to tumor necrosis factor-alpha. *J. Biol. Chem.*2005; 280:27879-87.

Membrane, caveola
{ECO:0000250|UniProtKB:Q9JHE3};
Single-pass type II membrane protein
{ECO:0000250|UniProtKB:Q91XT9}. Golgi
apparatus membrane; Single-pass type II
membrane protein
{ECO:0000250|UniProtKB:Q91XT9}.
Mitochondrion. Secreted, extracellular
exosome. Note=Enriched in exosomes upon
stimulation by cytokine
(PubMed:24798654). Enriched in caveolae
and lipid rafts (By similarity). The
localization to the mitochondrion could not
be confirmed (PubMed:15845354)
{ECO:0000250|UniProtKB:Q9JHE3,
ECO:0000269|PubMed:15845354,
ECO:0000269|PubMed:24798654}

Tissue Location

Primarily expressed in intestine
(PubMed:17334805). Ubiquitously
expressed with higher levels in kidney,
skeletal muscle and heart
(PubMed:10781606). The ubiquitous
expression observed for ASAH2 might be an
experimental artifact due to the paralog
ASAH2B (PubMed:17334805).

ASAH2 Antibody - Protocols

Provided below are standard protocols that you
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)