


**PALB2 Antibody**  
Catalog # ASC10949

**Specification**

**PALB2 Antibody - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">O86YC2</a>
Other Accession	<a href="#">O86YC2</a> , <a href="#">74727919</a>
Reactivity	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>IgG</b>
Calculated MW	<b>Predicted: 65, 100, 130 kDa</b>

Application Notes **Observed: 140 kDa**  
**PALB2 antibody can be used for detection of PALB2 by Western blot at 1 - 2 µg/mL.**



**PALB2 Antibody - Additional Information**

Gene ID **79728**  
**Target/Specificity**  
PALB2; At least four isoforms of PALB2 are known to exist; this antibody will detect all four isoforms.

**Reconstitution & Storage**  
PALB2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

**Precautions**  
PALB2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**PALB2 Antibody - Protein Information**

**Name** PALB2

**PALB2 Antibody - Background**

PALB2 Antibody: The breast cancer early onset proteins BRCA1 and BRCA2 are central to the repair of DNA damage by homologous recombination and are strongly associated with inherited breast and ovarian cancer. The protein PALB2 is required for the localization of BRCA2 to sites of DNA damage and like BRCA1 and 2 is a breast cancer susceptibility gene. PALB2 is thought to function through directly binding to BRCA1, which allows the PALB2 protein to organize BRCA2 and the recombinase RAD51 at the site of DNA damage. Analysis of the BRCA1-PALB2-BRCA2-RAD51 network may allow predictions to be made of the responsiveness of a particular tumor to therapeutic treatments.

**PALB2 Antibody - References**

Greenberg RA. Recognition of DNA double strand breaks by the BRCA1 tumor suppressor network. *Chromosoma* 2008; 117:305-17.  
Xia B, Sheng Q, Nakanishi K, et al. Control of

## Synonyms FANCN

### Function

Plays a critical role in homologous recombination repair (HRR) through its ability to recruit BRCA2 and RAD51 to DNA breaks (PubMed:<a href="http://www.uniprot.org/citations/16793542" target="\_blank">16793542</a>, PubMed:<a href="http://www.uniprot.org/citations/19423707" target="\_blank">19423707</a>, PubMed:<a href="http://www.uniprot.org/citations/19369211" target="\_blank">19369211</a>, PubMed:<a href="http://www.uniprot.org/citations/22941656" target="\_blank">22941656</a>, PubMed:<a href="http://www.uniprot.org/citations/24141787" target="\_blank">24141787</a>, PubMed:<a href="http://www.uniprot.org/citations/28319063" target="\_blank">28319063</a>). Strongly stimulates the DNA strand-invasion activity of RAD51, stabilizes the nucleoprotein filament against a disruptive BRC3-BRC4 polypeptide and helps RAD51 to overcome the suppressive effect of replication protein A (RPA) (PubMed:<a href="http://www.uniprot.org/citations/20871615" target="\_blank">20871615</a>). Functionally cooperates with RAD51AP1 in promoting of D-loop formation by RAD51 (PubMed:<a href="http://www.uniprot.org/citations/20871616" target="\_blank">20871616</a>). Serves as the molecular scaffold in the formation of the BRCA1-PALB2-BRCA2 complex which is essential for homologous recombination (PubMed:<a href="http://www.uniprot.org/citations/19369211" target="\_blank">19369211</a>). Via its WD repeats is proposed to scaffold a HR complex containing RAD51C and BRCA2 which is thought to play a role in HR-mediated DNA repair (PubMed:<a href="http://www.uniprot.org/citations/24141787" target="\_blank">24141787</a>). Essential partner of BRCA2 that promotes the localization and stability of BRCA2 (PubMed:<a href="http://www.uniprot.org/citations/16793542" target="\_blank">16793542</a>). Also enables its recombinational repair and checkpoint functions of BRCA2 (PubMed:<a href="http://www.uniprot.org/citations/1679

BRCA2 cellular and clinical functions by a nuclear partner, PALB. Mol. Cell 2007; 22:719-29.

Rahman N, Seal S, Thompson D, et al. PALB2, which encodes a BRCA2-interacting protein, is a breast cancer susceptibility gene. Nat. Genet. 2007; 39:165-7.

Zhang F, Fan Q, Ren K, et al. PALB2 functionally connects the breast cancer susceptibility proteins BRCA1 and BRCA2. Mol. Cancer. Res. 2009; 7:1110-8.

3542" target="\_blank">16793542</a>). May act by promoting stable association of BRCA2 with nuclear structures, allowing BRCA2 to escape the effects of proteasome-mediated degradation (PubMed:<a href="http://www.uniprot.org/citations/16793542" target="\_blank">16793542</a>). Binds DNA with high affinity for D loop, which comprises single-stranded, double-stranded and branched DNA structures (PubMed:<a href="http://www.uniprot.org/citations/20871616" target="\_blank">20871616</a>). May play a role in the extension step after strand invasion at replication-dependent DNA double-strand breaks; together with BRCA2 is involved in both POLH localization at collapsed replication forks and DNA polymerization activity (PubMed:<a href="http://www.uniprot.org/citations/24485656" target="\_blank">24485656</a>).

#### Cellular Location

Nucleus Note=Colocalizes with BRCA2 and BRCA1 in nuclear foci

#### PALB2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)