

OCIAD2 Antibody
Catalog # ASC11007

Specification

OCIAD2 Antibody - Product Information

Application	WB, IF
Primary Accession	Q56VL3
Other Accession	NP_001014446 , 62244044
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	OCIAD2 antibody can be used for detection of OCIAD2 by Western blot at 0.5 - 1 µg/mL. Antibody can also be used for immunofluorescence starting at 5 µg/mL. For immunofluorescence start at 5 µg/mL.

OCIAD2 Antibody - Additional Information

Gene ID	132299
Target/Specificity	OCIAD2;

Reconstitution & Storage


OCIAD2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

OCIAD2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

OCIAD2 Antibody - Protein Information

Name OCIAD2



Western blot analysis of OCIAD2 in A549 cell lysate with OCIAD2 antibody at (A) 0.5 and (B) 1 µg/mL.

Immunofluorescence of OCIAD2 in A549 cells with OCIAD2 antibody at 5 µg/mL.

OCIAD2 Antibody - Background

OCIAD2 Antibody: OCIAD2 was identified by its sequence similarity with OCIAD1, and together OCIAD1 and OCIAD2 form the OCIA domain family. OCIAD2 mRNA was found to be expressed at higher levels in invasive adenocarcinoma mixed subtype with bronchioloalveolar carcinoma component (BAC) of the lung. Loss of OCIAD2 expression was significantly correlated with lymphatic invasion, blood vessel invasion, and lymph node metastasis, indicating that OCIAD2 may

Cellular Location
Endosome.

OCIAD2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

play a role in cell adhesion and prevention of cell migration. While the function of OCIAD2 is still unknown, its expression in adenocarcinoma with BAC component is significantly associated with a favorable prognosis and may serve as a marker for selecting tumors that are treatable by limited surgery.

OCIAD2 Antibody - References

Strausberg RL, Feingold EA, Grouse LH, et al. Mammalian gene collection program team. Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. Proc. Natl. Acad. Sci. USA 2002; 99:16899-903.

Ishiyama T, Kano J, Anami Y, et al. OCIA domain containing 2 is highly expressed in adenocarcinoma mixed subtype with bronchioloalveolar carcinoma component and is associated with better prognosis. Cancer Sci. 2006; 98:50-7.