

MINA Antibody
Catalog # ASC11072

Specification

MINA Antibody - Product Information

Application	WB, IHC, IF
Primary Accession	Q8IUJ8
Other Accession	NP_694822 , 110227621
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	MINA antibody can be used for detection of MINA by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.

MINA Antibody - Additional Information

Gene ID **84864**
Target/Specificity
MINA;

Reconstitution & Storage


MINA antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions


MINA Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

MINA Antibody - Protein Information

Name RIOX2 ([HGNC:19441](#))



Western blot analysis of MINA in human heart tissue lysate with MINA antibody at (A) 1 and (B) 2 µg/mL.



Immunohistochemistry of MINA in mouse heart tissue with MINA antibody at 5 µg/mL.

Function

Oxygenase that can act as both a histone lysine demethylase and a ribosomal histidine hydroxylase. Is involved in the demethylation of trimethylated 'Lys-9' on histone H3 (H3K9me3), leading to an increase in ribosomal RNA expression. Also catalyzes the hydroxylation of 60S ribosomal protein L27a on 'His-39'. May play an important role in cell growth and survival. May be involved in ribosome biogenesis, most likely during the assembly process of pre-ribosomal particles.

Cellular Location

Nucleus. Nucleus, nucleolus


Tissue Location

Expressed in liver, skeletal muscle, heart, pancreas, and placenta. Not detected in brain, lung or kidney. Expressed in several lung cancer tissues, but is barely detected in the adjacent non-cancerous tissues. Also highly expressed in several esophageal squamous cell carcinoma (ESCC), and colon cancer tissues, and in various cancer cell lines.

MINA Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Immunofluorescence of MINA in mouse heart tissue with MINA antibody at 20 µg/mL.

MINA Antibody - Background

MINA Antibody: MINA is nuclear localized, myc-inducible protein that is thought to play a role in mammalian cell proliferation. Treatment of cancer cell lines such as the colon cancer cell line SW680 with siRNA against MINA inhibits cell growth, demonstrating that MINA may be a potential therapeutic target. MINA regulates several genes related to cell adhesion and metabolism that have also been shown to be regulated by c-Myc, but also regulates other genes whose expression are not modulated by c-Myc such as EGFR, IL-6 and HGF. MINA has also been found to act as a repressor to IL-4 expression in T cells, indicating that it may also play a role in T cell differentiation and genetic variation in T helper type 2 bias.

MINA Antibody - References

- Tsuneoka M, Kody Y, Soejima M, et al. A novel myc target gene, mina53, that is involved in cell proliferation. *J. Biol. Chem.*2002; 277:35450-9.
- Teye K, Tsuneoka M, Arima N, et al. Increased expression of a Myc target gene Mina53 in human colon cancer. *Am. J. Pathol.*2004; 164:205-16.
- Komiya K, Sueoka-Aragane N, Sato A, et al. Mina53, a novel c-Myc target gene, is frequently expressed in lung cancers and exerts oncogenic property in NIH/3T3 cells. *J. Cancer Res. Clin. Oncol.*2010; 136:465-73.
- Okamoto M, Van Stry M, Chung L, et al. Mina, an IL4 repressor, controls T helper type 2 bias.

Nat. Immunol.2009; 10:872-9.