

## DBX1 Antibody Catalog # ASC11366

# **Specification**

### **DBX1 Antibody - Product Information**

Application Primary Accession Other Accession

A6NMT0 NP\_001025036, 304434796

WB, IHC, IF

Reactivity

Human, Mouse, Rat

Host Clonality Isotype Rabbit Polyclonal IgG

Application Notes

DBX1 antibody can be used for detection of DBX1 by Western blot at 1 - 2 µg/mL. Antibody can also be used for immu nohistochemistry starting at 2.5 µg/mL. For immun of luorescence

ofluorescence start at 20 μg/mL.

# **DBX1** Antibody - Additional Information

Gene ID **120237** 

**Target/Specificity** 

DBX1; BX1 antibody is predicted to not cross-react with other DBX protein family members


### **Reconstitution & Storage**

DBX1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.


## **Precautions**

DBX1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**DBX1 Antibody - Protein Information** 



Western blot analysis of DBX1 in mouse kidney tissue lysate with DBX1 antibody at (A) 1 and (B) 2  $\mu$ g/mL.



Immunohistochemistry of DBX1 in human kidney tissue with DBX1 antibody at 2.5  $\mu$ g/mL.



#### Name DBX1

#### **Function**

Could have a role in patterning the central nervous system during embryogenesis. Has a key role in regulating the distinct phenotypic features that distinguish two major classes of ventral interneurons, V0 and V1 neurons. Regulates the transcription factor profile, neurotransmitter phenotype, intraspinal migratory path and axonal trajectory of V0 neurons, features that differentiate them from an adjacent set of V1 neurons (By similarity).


### **Cellular Location**

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108}.

# **DBX1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture



Immunofluorescence of DBX1 in human kidney tissue with DBX1 antibody at 20 µg/mL.

### **DBX1 Antibody - Background**

DBX1 Antibody: DBX1 is implicated in patterning the central nervous system during embryogenesis. DBX1 homeodomain transcription factor is expressed in progenitors which are critical in establishing cell fate allocation and cell diversity. DBX1 has a key role in regulating the distinct phenotypic features that distinguish two major classes of ventral interneurons, V0 and V1 neurons. DBX1-derived neurons are essential for the transgenic expression and generation of respiratory behavior both in vitro and in vivo.

# **DBX1 Antibody - References**

Griveau A, Borello U, Causeret F, et al. A novel role for Dbx1-derived Cajal-Retzius cells in early regionalization of the cerebral cortical neuroepithelium. PLoS Biol. 2010; 8(7):e1000440

Causeret F, Ensini M, Teissier A, et al. Dbx1-expressing cells are necessary for the survival of the Mammalian anterior neural and craniofacial structures. PLoS One 2011; 6:e19367.

Pierani A, Moran-Rivard L, Sunshine MJ, et al. Control of interneuron fate in the developing spinal cord by the progenitor homeodomain protein Dbx1. Neuron 2001; 29:367-84 Gray PA, Hayes JA, Ling GY, et al. Developmental origin of preBotzinger complex respiratory neurons. J. Neurosci. 2010; 30:14883-95.