

**Alginate Antibody**  
**Alginate Antibody, Clone 4B10-1C5**  
**Catalog # ASM10153**

**Specification**

**Alginate Antibody - Product Information**

Application **IHC, WB**  
Host **Mouse**  
Isotype **IgG1 Kappa**  
Clonality **Monoclonal**

**Description**

Mouse Anti- Alginate Monoclonal IgG1  
Kappa

**Target/Specificity**

Binds selectively to a BSA-conjugated alginate, but not to unconjugated BSA.

**Other Names**

Alginic Acid Antibody, Algin Antibody,  
Sodium Alginate Antibody

**Immunogen**

Sodium Alginate conjugated to KLH

**Purification**

Protein G Purified

Storage **-20°C**

**Storage Buffer**

PBS pH7.4, 50% glycerol, 0.09% sodium  
azide

Shipping **Blue Ice or 4°C**  
Temperature


**Certificate of Analysis**

1 µg/ml of SMC-208 was sufficient for  
detection of 150 ng of alginate-conjugated  
BSA by colorimetric immunoblot analysis  
using Goat anti-mouse IgG:HRP as the  
secondary antibody.


**Alginate Antibody - Protocols**

Provided below are standard protocols that you  
may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)



Immunohistochemistry analysis using Mouse Anti-Alginate Monoclonal Antibody, Clone 4B10-1C5 (ASM10153). Tissue: Burned Skin. Species: Human. Primary Antibody: Mouse Anti-Alginate Monoclonal Antibody (ASM10153) at 1:200.



Western Blot analysis of ALL BSA-Alginate Conjugate showing detection of ~250 kDa Alginate protein using Mouse Anti-Alginate Monoclonal Antibody, Clone 4B10-1C5 (ASM10153). Lane 1: MW ladder. Lane 2: 0.625ug BSA:Alginate. Load: 0.625 µg. Block: 5% milk + TBST for 1 hour at RT. Primary Antibody: Mouse Anti-Alginate Monoclonal

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Antibody (ASM10153) at 1:500 for 1 hour at RT. Secondary Antibody: HRP Goat Anti-Mouse at 1:100 for 1 hour at RT. Color Development: TMB solution for 2 min at RT. Predicted/Observed Size: ~250 kDa.

### **Alginate Antibody - Background**

Sodium alginate is used in biological experiments for the immobilization of cells and encapsulation due to its biocompatibility and simple gelation with divalent cations such as  $\text{Ca}^{2+}$ . Studies suggest that preparation of alginate microspheres will sustain protein delivery within tissue scaffolds (1), as well as in many other cell types (2).

### **Alginate Antibody - References**

1. Zhai P., Chen X.B., and Schreyer D.J. (2013) Biofabrication. 5(1): 015009.
2. Selimoglu S.M., Ayyildiz-Tamis D., Gurhan I.D., and Elibol M. (2012) J Biosci Bioeng. 113(2):233-238.