

**N-terminal Arginylation Antibody**  
**N-terminal Arginylation Antibody, Clone 2A4**  
**Catalog # ASM10170**

**Specification**

**N-terminal Arginylation Antibody - Product Information**

Application **WB**  
Host **Mouse**  
Isotype **IgG3**  
Clonality **Monoclonal**

**Description**

Mouse Anti-N-terminal Arginylation Monoclonal IgG3

**Target/Specificity**

Specific for N-terminal arginine, next to both glutamic acid and aspartic acid. Does not detect internal arginine.

**Other Names**

N-terminal Arginine Antibody, N-terminal Arginylation Antibody, N-terminal Arginylated Antibody, N terminal Arginine Antibody, N terminal Arginylation Antibody, N terminal Arginylated Antibody, Amino-terminal Arginine Antibody, Amino-terminal Arginylation Antibody, Amino-terminal Arginylated Antibody, Amino terminal Arginine Antibody, Amino terminal Arginylation Antibody, Amino terminal Arginylated Antibody

**Immunogen**

Synthetic N-terminal arginylated peptide conjugated to KLH

**Purification**

Protein G Purified

Storage **-20°C**

**Storage Buffer**


PBS pH 7.4, 50% glycerol, 0.9% Sodium Azide

Shipping **Blue Ice or 4°C**

Temperature

**Certificate of Analysis**

A 1:1000 dilution of SMC-262 was sufficient for detection of N-terminal Arginylation in 0.5 ug of N-terminal Arginine peptide conjugated to BSA by ECL immunoblot



**N-terminal Arginylation Antibody - Background**

Protein arginylation is the post-translational addition of arginine to proteins by arginyltransferase ATE1. Arginylation of proteins has been found to play an essential role in physiological pathways during embryogenesis and adulthood (1). Arginylation

analysis using goat anti-mouse IgG:HRP as the secondary antibody.

**Cellular Localization**  
Endoplasmic Reticulum

**N-terminal Arginylation Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

has also been shown to regulate cell stress responses, including ER stress, cytosolic misfolded proteins, and heat stress (2).

**N-terminal Arginylation Antibody - References**

1. Saha S. and Kashina A. (2011) Dev Biol. 385(1): 1-8.
2. Deka K., et al. (2016) Cell Death Discov. 2: 16074.