

**KCNQ4 Antibody**  
**KCNQ4 Antibody, Clone S43-6**  
**Catalog # ASM10187**

**Specification**

**KCNQ4 Antibody - Product Information**

Application **IHC, WB**  
 Primary Accession [P56696](#)  
 Other Accession [NP\\_004691.2](#)  
 Host **Mouse**  
 Isotype **IgG1**  
 Reactivity **Human, Mouse, Rat**  
 Clonality **Monoclonal**

**Description**

Mouse Anti-Human KCNQ4 Monoclonal IgG1

**Target/Specificity**

Detects ~77kDa.

**Other Names**

DFNA 2 antibody, DFNA2 antibody, KCNQ 4 antibody, Kcnq4 antibody, KCNQ4\_HUMAN antibody, KQT like 4 antibody, KQT-like 4 antibody, KV7.4 antibody, Potassium channel KQT like 4 antibody, Potassium channel subunit alpha KvLQT4 antibody, Potassium voltage gated channel KQT like protein 4 antibody, Potassium voltage gated channel KQT like subfamily member 4 antibody, Potassium voltage gated channel subfamily KQT member 4 antibody, Potassium voltage-gated channel subfamily KQT member 4 antibody, Voltage gated potassium channel subunit Kv7.4 antibody, Voltage-gated potassium channel subunit Kv7.4 antibody

**Immunogen**

Fusion protein amino acids 2-77 of human KCNQ4

**Purification**


Protein G Purified

Storage **-20°C**


**Storage Buffer**

PBS pH7.4, 50% glycerol, 0.09% sodium azide

Shipping **Blue Ice or 4°C**  
 Temperature



Immunohistochemistry analysis using Mouse Anti-KCNQ4 Monoclonal Antibody, Clone S43-6 (ASM10187). Tissue: frozen brain section. Species: mouse. Fixation: 10% Formalin Solution for 12-24 hours at RT. Primary Antibody: Mouse Anti-KCNQ4 Monoclonal Antibody (ASM10187) at 1:1000 for 1 hour at RT. Secondary Antibody: HRP/DAB Detection System: Biotinylated Goat Anti-Mouse, Streptavidin Peroxidase, DAB Chromogen (brown) for 30 minutes at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 250-500 µl for 5 minutes at RT.



Western Blot analysis of Rat tissue lysate

### Certificate of Analysis

1 µg/ml of SMC-309 was sufficient for detection of KCNQ4 in 10 µg of COS-1 cell lysate transiently expressing KCNQ4 by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

### Cellular Localization


Plasma Membrane | Basal Cell Membrane

### KCNQ4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

showing detection of KCNQ4 protein using Mouse Anti-KCNQ4 Monoclonal Antibody, Clone S43-6 (ASM10187). Load: 15 µg. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-KCNQ4 Monoclonal Antibody (ASM10187) at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.



Immunohistochemistry analysis using Mouse Anti-KCNQ4 Monoclonal Antibody, Clone S43-6 (ASM10187). Tissue: hippocampus. Species: Human. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-KCNQ4 Monoclonal Antibody (ASM10187) at 1:1000 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT.

### KCNQ4 Antibody - Background

The protein encoded by this gene forms a potassium channel that is thought to play a critical role in the regulation of neuronal excitability (1), particularly in sensory cells of the cochlea (2). The current generated by this channel is inhibited by M1 muscarinic acetylcholine receptors and activated by retigabine, a novel anti-convulsant drug (3).

### KCNQ4 Antibody - References

1. Hernandez C.C., Zaiko O., Tolstykh G.P., Shapiro M.S. (2008) J Physiol. 586(7): 1811-1821.
2. Kharkovets T., et al. (2006) EMBO J. 25(3): 642-652.
3. Tatulian L., Delmas P., Abogadie F.C., Brown D.A. (2001) J Neuroscience. 21(15): 5535-5545.