

**SHANK1 Antibody**  
**SHANK1 Antibody, Clone S22-21**  
**Catalog # ASM10205**

**Specification**

**SHANK1 Antibody - Product Information**

Application **IHC, WB**  
Primary Accession [O9WV48](#)  
Other Accession [NP\\_113939.2](#)  
Host **Mouse**  
Isotype **IgG1**  
Reactivity **Human, Mouse, Rat**  
Clonality **Monoclonal**

**Description**

Mouse Anti-Rat SHANK1 Monoclonal IgG1

**Target/Specificity**

Detects ~190-220 kDa (alternative splice variants). No cross-reactivity against Shank2 or Shank3.

**Other Names**

GKAP/SAPAP interacting protein Antibody, OTTHUMP00000174437 Antibody, SH3 and multiple ankyrin repeat domains 1 Antibody, SH3 and multiple ankyrin repeat domains protein 1 Antibody, SH3/ankyrin domain gene 1 Antibody, SHAN1\_HUMAN Antibody, SHANK 1 Antibody, Shank1 Antibody, Shank1a Antibody, Somatostatin receptor interacting protein Antibody, Somatostatin receptor-interacting protein Antibody, SPANK 1 Antibody, SPANK1 Antibody, SSTR interacting protein Antibody, SSTR-interacting protein Antibody, SSTRIP Antibody, Synamon Antibody

**Immunogen**

Fusion protein amino acids 469-691 (SH3/PDZ domains) of rat Shank1

**Purification**


Protein G Purified

Storage **-20°C**


**Storage Buffer**

PBS pH7.4, 50% glycerol, 0.09% sodium azide

Shipping **Blue Ice or 4°C**



Immunohistochemistry analysis using Mouse Anti-SHANK1 Monoclonal Antibody, Clone S22-21 (ASM10205). Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-SHANK1 Monoclonal Antibody (ASM10205) at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Filaggrin-like staining (upper layer aggregations of staining).



Western Blot analysis of Rat brain membrane lysate showing detection of SHANK1 protein using Mouse Anti-SHANK1 Monoclonal Antibody, Clone S22-21

Temperature

### Certificate of Analysis

1 µg/ml of SMC-329 was sufficient for detection of Shank1 in 10 µg of rat brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

### Cellular Localization

Cytoplasm | Cell Junction | Synapse | Postsynaptic Cell Membrane | Postsynaptic Density

### SHANK1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

(ASM10205). Load: 15 µg. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-SHANK1 Monoclonal Antibody (ASM10205) at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.

### SHANK1 Antibody - Background

Shank proteins make up a family of scaffold proteins identified through their interaction with a variety of membrane and cytoplasmic proteins (1). Shank proteins at postsynaptic sites of excitatory synapses play roles in signal transmission into the postsynaptic neuron. Studies suggest that Shank2 is expressed in the neurons of the developing retina, and could play a role in the neuronal differentiation of the developing retina (2). Other recent studies suggest that the disruption of glutamate receptors at the Shank postsynaptic platform could contribute to the destruction of the postsynaptic density, which underlies the synaptic dysfunction and loss in Alzheimer's disease (3).

### SHANK1 Antibody - References

1. Sheng M., and Kim E. (2000) Journal of Cell Science. 113: 1851-1856.
2. Kim J.H., et al. (2009) Exp Mol Med. 41(4): 236-242.
3. Gong Y., et al. (2009) Brain Res. 1292: 191-198.