

GABA-A Receptor Beta1 Antibody

GABA A Receptor Beta1 Antibody, Clone S96-55 Catalog # ASM10216

Specification

GABA-A Receptor Beta1 Antibody - Product Information

Application	ICC/IF, WB, IHC
Primary Accession	<u>P50571</u>
Other Accession	<u>NP_032095.1</u>
Host	Mouse
lsotype	lgG1
Reactivity	Human, Mouse,
	Rat
Clonality	Monoclonal

Description Mouse Anti-Mouse GABA-A Receptor Beta1 Monoclonal IgG1

HRP

Target/Specificity Detects ~55kDa. No cross-reactivity against GABA-A-R-Beta 2 or -Beta3.

Other Names

Format


AW061132 antibody, B230208N19Rik antibody, GABA(A) receptor beta 1 antibody, GABA(A) receptor subunit beta-1 antibody, GABA-A receptor beta-1 polypeptide antibody, Gabrb-1 antibody, GABRB1 antibody, Gamma aminobutyric acid (GABA) A receptor beta 1 antibody, Gamma Aminobutyric Acid A Receptor Beta 1 antibody, Gamma Aminobutyric Acid Receptor beta-1 antibody, Gamma-aminobutyric acid (GABA) A receptor subunit beta 1 antibody, Gamma-aminobutyric acid receptor subunit beta-1 antibody, GARB1 antibody, GBRB1_HUMAN antibody

Immunogen

Fusion protein amino acids 327-450 of mouse GABA-A-R-Beta1

Purification Protein G Purified

Storage -20°C Storage Buffer PBS pH7.4, 50% glycerol, 0.09% sodium azide



Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-GABA A Receptor Monoclonal Antibody, Clone S96-55 (ASM10216). Tissue: HaCaT cells. Species: Human. Fixation: Cold 100% methanol for 10 minutes at -20°C. Primary Antibody: Mouse Anti-GABA A Receptor Monoclonal Antibody (ASM10216) at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Positive.

201.5→
156.75→ 106→
79.68→
48.33→
37.81→
23.27→
18.19→

Western Blot analysis of Human Cell line lysates showing detection of GABA A Receptor protein using Mouse Anti-GABA A Receptor Monoclonal Antibody, Clone S96-55



Shipping

Blue Ice or 4ºC

Temperature **Certificate of Analysis** 2 μ g/ml of SMC-340 was sufficient for detection of Beta1 GABA receptor in 10 μ g of rat brain lysate by colorimetric immunoblot analysis using goat anti-mouse lgG:HRP as the secondary antibody.


Cellular Localization Cell Membrane | Cell Junction | Synapse | Postsynaptic Cell Membrane

GABA-A Receptor Beta1 Antibody -Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

(ASM10216). Load: 15 µg. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-GABA A Receptor Monoclonal Antibody (ASM10216) at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.



Immunohistochemistry analysis using Mouse Anti-GABA A Receptor Monoclonal Antibody, Clone S96-55 (ASM10216). Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-GABA A Receptor Monoclonal Antibody (ASM10216) at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Dermal staining.

GABA-A Receptor Beta1 Antibody -Background

The GABA-A receptor is a member of the superfamily of fast acting ligand-gated ion channels. The individual subunits of these receptors have similar sequences and structural features (1). GABA-A receptors are the major fast inhibitory neurotransmitter gated ion channels in the brain (2).

GABA-A Receptor Beta1 Antibody -References

 Bracamontes J.R. and Steinbach J.H. (2008)
J Bio Chem. 283: 26128-26136.
Macdonald R.L., Olsen R.W. (1993) Annu Rev Neurosci. 17: 569-602.