

LGI1 Antibody
LGI1 Antibody, Clone S283-7
Catalog # ASM10293

Specification

LGI1 Antibody - Product Information

Application	ICC/IF, WB
Primary Accession	O9JIA1.1
Other Accession	NP_064674.1
Host	Mouse
Isotype	IgG2a
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Format	HRP

Description

Mouse Anti-Mouse LGI1 Monoclonal IgG2a

Target/Specificity

Detects ~60kDa.

Other Names

ADLTE Antibody, ADPAEF Antibody, ADPEAF Antibody, Epitempin 1 Antibody, EPT Antibody, ETL1 Antibody, IB1099 Antibody, leucine rich glioma inactivated 1 Antibody, OTTHUMP00000020121 Antibody, OTTHUMP00000020122 Antibody

Immunogen

Fusion protein amino acids 37-113 (LRRNT domain and first LRR repeat) of mouse LGI1. Rat: 100% identity (77/77 amino acids identical). Human: 98% identity (76/77 amino acids identical). ~50% identity with LGI2, LGI3 and LGI4.

Purification

Protein G Purified

Storage **-20°C**


Storage Buffer

PBS pH 7.4, 50% glycerol, 0.1% sodium azide

Shipping **Blue Ice or 4°C**
Temperature

Certificate of Analysis

1 µg/ml of SMC-461 was sufficient for detection of LGI1 in 20 µg of rat brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary



Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-LGI1 Monoclonal Antibody, Clone S283-7 (ASM10293). Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-LGI1 Monoclonal Antibody (ASM10293) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Cell Junction, Golgi Apparatus, Endoplasmic Reticulum. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) LGI1 Antibody (D) Composite.

antibody.


Cellular Localization

Cell Junction | Golgi Apparatus |
Endoplasmic Reticulum

LG11 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



LG11 Antibody - Background

The leucine-rich, glioma inactivated gene 1 (LG1) was first identified as a candidate tumor suppressor gene for glioma and may play a role in other cancers. LG1 is a member of a family of highly related proteins containing leucine-rich repeats (LRRs) which are highly similar to other transmembrane signaling molecules and receptors. LG1 serves as a ligand to ADAM22, a metalloprotease localized at the synapse. Mutations in LG1 account for nearly half of autosomal dominant lateral temporal epilepsy (ADTLE), an epileptic syndrome characterized by focal seizures with predominant auditory symptoms. Two isoforms of LG1 are known to exist; this LG1 antibody will recognize only the longer form.

LG11 Antibody - References

1. Chernova O.B., Somerville R.P. and Cowell J.K. (1998) *Oncogene*. 17:2873-81.
2. Fialka F., et al. (2008) *Oral Oncol*.
3. Gu W., et al. (2005) *Mol. Biol. Evol.* 22:2209-16.
4. Fukata Y., et al. (2006) *Science*. 313:1792-5.

