

REEP2 Antibody
REEP2 Antibody, Clone S326D-2
Catalog # ASM10313

Specification

REEP2 Antibody - Product Information

Application	ICC/IF, WB
Primary Accession	O8VCD6
Other Accession	NP_001191843.1
Host	Mouse
Isotype	IgG2A
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Format	RPE

Description
Mouse Anti-Mouse REEP2 Monoclonal IgG2A

Target/Specificity
Detects ~30kDa.

Other Names
Receptor expression-enhancing protein 2 Antibody, C2orf19 Antibody, SGC32445 Antibody, LOC682105 Antibody

Immunogen
Fusion protein amino acids 111-254 (cytoplasmic C-terminus) of mouse REEP2


Purification
Protein G Purified

Storage **-20°C**
Storage Buffer
PBS pH7.4, 50% glycerol, 0.1% sodium azide

Shipping **Blue Ice or 4°C**
Temperature

Certificate of Analysis
A 1:100 dilution of SMC-481 was sufficient for detection of REEP2 in 20 µg of mouse brain lysate by ECL immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

Cellular Localization
Mitochondrion | Mitochondrion Membrane | Endoplasmic Reticulum | Endoplasmic Reticulum Membrane




Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-REEP2 Monoclonal Antibody, Clone S326D-2 (ASM10313). Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-REEP2 Monoclonal Antibody (ASM10313) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000; 1:5000 for 60 min RT, 5 min RT. Localization: Mitochondrion, Mitochondrion Membrane, Endoplasmic Reticulum, Endoplasmic Reticulum Membrane. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) REEP2 Antibody (D) Composite.

REEP2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



REEP2 Antibody - Background

REEP2 belongs to the REEP family, which are transmembrane proteins that interact with odorant receptor proteins and may enhance the odorant receptor responses to odorants. REEP2 is also an integral membrane protein expressed in taste cells. Unlike its counterpart REEP1, REEP2 does not increase cell surface expression to sweet receptors, but instead alters their spatial organization (1).

REEP2 Antibody - References

1. Ilegems E., et al. (2010) J Neurosci. 30(41): 13774-13783.