

HSP90 Antibody
Catalog # **ASM10358**

Specification

HSP90 Antibody - Product Information

Application **ICC/IF, WB, IHC**
 Primary Accession [P08238](#)
 Other Accession [NP_031381.2](#)
 Host **Rabbit**
 Reactivity **Human, Mouse, Rat**
 Clonality **Polyclonal**

Description
 Rabbit Anti-Human HSP90 Polyclonal

Target/Specificity
 Detects ~90kDa.

Other Names
 HSP84 Antibody, HSP86 Antibody, HSP90A Antibody, HSP90AA1 Antibody, HSP90AB1 Antibody, HSP90B Antibody, HSPC1 Antibody, HSPC2 Antibody, HSPCAL1 Antibody, HSPCAL4 Antibody, HSP90N Antibody

Immunogen
 Full length protein HSP90

Purification
 Protein A Purified

Storage **-20°C**
Storage Buffer
 PBS pH 7.4, 50% glycerol, 0.09% sodium azide


Shipping **Blue Ice or 4°C**
 Temperature

Certificate of Analysis
 A 1:500 dilution of SPC-104 was sufficient for detection of 0.2 mg of purified HSP90 by ECL immunoblot analysis.


Cellular Localization
 Cytoplasm | Melanosome

HSP90 Antibody - Protocols

Provided below are standard protocols that you




Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-Hsp90 Polyclonal Antibody (ASM10358). Tissue: Heat Shocked HeLa Cells. Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Rabbit Anti-Hsp90 Polyclonal Antibody (ASM10358) at 1:100 for 12 hours at 4°C. Secondary Antibody: FITC Goat Anti-Rabbit (green) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Cytoplasm. Magnification: 100x. (A) DAPI (blue) nuclear stain. (B) Anti-Hsp90 Antibody. (C) Composite. Heat Shocked at 42°C for 1h.




Western blot analysis of Human Cell line lysates showing detection of HSP90 protein using Rabbit Anti-HSP90 Polyclonal Antibody (ASM10358). Load: 2 µg Hsp90beta. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Rabbit Anti-HSP90 Polyclonal Antibody (ASM10358) at 1:10000 for 2 hours at RT. Secondary Antibody: Donkey Anti-Rabbit IgG: HRP for 1 hour at RT.

may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Immunohistochemistry analysis using Rabbit Anti-HSP90 Polyclonal Antibody (ASM10358). Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative Solution. Primary Antibody: Rabbit Anti-HSP90 Polyclonal Antibody (ASM10358) at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Rabbit (green) at 1:50 for 1 hour at RT. Localization: Cytoplasm.



Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-Hsp90 Polyclonal Antibody (ASM10358). Tissue: Heat Shocked HeLa Cells. Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Rabbit Anti-Hsp90 Polyclonal Antibody (ASM10358) at 1:100 for 12 hours at 4°C. Secondary Antibody: APC Goat Anti-Rabbit (red) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Cytoplasm. Magnification: 20x. (A) DAPI (blue) nuclear stain. (B) Anti-Hsp90 Antibody. (C) Composite. Heat Shocked at 42°C for 1h.

HSP90 Antibody - Background

HSP90 is a highly conserved and essential stress protein that is expressed in all eukaryotic cells. From a functional perspective, HSP90 participates in the folding, assembly, maturation, and stabilization of specific proteins as an integral component of a chaperone complex (1-4). Despite its label of being a heat-shock protein, HSP90 is one of the most highly expressed proteins in unstressed cells (1-2% of cytosolic protein). It carries out a number of housekeeping functions - including

controlling the activity, turnover, and trafficking of a variety of proteins. Most of the HSP90-regulated proteins that have been discovered to date are involved in cell signaling (5-6). The number of proteins now known to interact with HSP90 is about 100. Target proteins include the kinases v-Src, Wee1, and c-Raf, transcriptional regulators such as p53 and steroid receptors, and the polymerases of the hepatitis B virus and telomerase.5. When bound to ATP, HSP90 interacts with co-chaperones Cdc37, p23, and an assortment of immunophilin-like proteins, forming a complex that stabilizes and protects target proteins from proteasomal degradation. In most cases, HSP90-interacting proteins have been shown to co-precipitate with HSP90 when carrying out immunoadsorption studies, and to exist in cytosolic heterocomplexes with it. In a number of cases, variations in HSP90 expression or HSP90 mutation has been shown to degrade signaling function via the protein or to impair a specific function of the protein (such as steroid binding, kinase activity) in vivo. Ansamycin antibiotics, such as geldanamycin and radicicol, inhibit HSP90 function (7). Looking for more information on HSP90? Visit our new HSP90 Scientific Resource Guide at <http://www.HSP90.ca>.

HSP90 Antibody - References

1. Arlander S.J.H., et al. (2003) J Biol Chem 278: 52572-52577.
2. Pearl H., et al. (2001) Adv Protein Chem 59:157-186.
3. Neckers L., et al. (2002) Trends Mol Med 8:S55-S61.
4. Pratt W., Toft D. (2003) Exp Biol Med 228:111-133.
5. Pratt W., Toft D. (1997) Endocr Rev 18: 306-360.
6. Pratt W.B. (1998) Proc Soc Exptl Biol Med 217: 420-434.
7. Whitesell L., et al. (1994) Proc Natl Acad Sci USA 91: 8324-8328.