

BIM/BOD (IN) Antibody
Catalog # ASM10372

Specification

BIM/BOD (IN) Antibody - Product Information

Application **IHC, WB**
 Primary Accession [O43521](#)
 Other Accession [NP_619527.1](#)
 Host **Rabbit**
 Reactivity **Human, Mouse, Rat**
 Clonality **Polyclonal**
Description
 Rabbit Anti-Human BIM/BOD (IN) Polyclonal

Target/Specificity
 Detects ~23kDa.

Other Names

BCL2 like protein 11 Antibody, Bcl2 interacting mediator of cell death Antibody, Bcl2 interacting protein Bim Antibody, BCL2 like 11 Antibody, BAM Antibody, BimE1 Antibody, BimL Antibody, BOD Antibody, BIM beta 6 Antibody, BIM Beta7 Antibody, BIM alpha 3 Antibody, BIM alpha4 Antibody, BIM alpha5 Antibody, BIM alpha6 Antibody

Immunogen

Internal central amino acids of human Bim

Purification

Protein A Purified

Storage **-20°C**

Storage Buffer

PBS, 50% glycerol, 0.09% sodium azide


Shipping **Blue Ice or 4°C**
 Temperature

Certificate of Analysis


1 µg/ml of SPC-113 was sufficient for detection of Bim in 20 µg of K562 and A549 cell lysates by colorimetric immunoblot analysis using Goat anti-rabbit IgG:HRP as the secondary antibody.

Cellular Localization

Mitochondrion | Endomembrane System



Immunohistochemistry analysis using Rabbit Anti-BIM Polyclonal Antibody (ASM10372). Tissue: skin cancer cells. Species: Human. Primary Antibody: Rabbit Anti-BIM Polyclonal Antibody (ASM10372) at 1:100.



Western blot analysis of Human K562 (Left) and A549 (Right) whole cell lysates showing detection of BIM protein using Rabbit Anti-BIM Polyclonal Antibody (ASM10372). Primary Antibody: Rabbit Anti-BIM Polyclonal Antibody (ASM10372) at 1:1000.

BIM/BOD (IN) Antibody - Background

Members in the Bcl-2 family are critical regulators of apoptosis by either inhibiting or promoting cell death. Bim/BOD is a group of three splice variants, BimEL, BimL and BimS,

BIM/BOD (IN) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

with apparent molecular masses of ~23, 16, and 13 kDa, respectively. Bcl-2 homology 3 (BH3) domain is a potent death domain. BH3 domain containing pro-apoptotic proteins, including Bad, Bax, Bid, Bik, and Hrk, form a growing subclass of the Bcl-2 family. A novel BH3 domain containing protein was recently identified and designated Bim or BOD in human, mouse and rat (1,2). Bim/BOD interacts with diverse members in the pro-survival Bcl-2 sub-family including Bcl-2, Bcl-xL and Bcl-w. Bim/BOD induces apoptosis. The messenger RNA of Bim is ubiquitously expressed in multiple tissues and cell lines (1,2).

BIM/BOD (IN) Antibody - References

1. O'Connor L., Strasser A., O'Reilly L.A., et al. (1998) EMBO J. 17: 384-395.
2. Hsu S.Y., Lin P., and Hsueh A.J. (1998) Mol. Endocrinol. 12: 1432-40.