

ATG4C Antibody
Catalog # ASM10514

Specification

ATG4C Antibody - Product Information

Application **ICC/IF, WB**
Primary Accession [Q96DT6](#)
Other Accession [NP_116241.2](#)
Host **Rabbit**
Reactivity **Human**
Clonality **Polyclonal**

Description

Rabbit Anti-Human ATG4C Polyclonal

Target/Specificity

Detects ~50 kDa.


Other Names

AUT like 1cysteine endopeptidase (*S. cerevisiae*) Antibody, APG4 autophagy 4 homolog C Antibody, Cysteine protease ATG4C Antibody, APG4-C Antibody, Autophagy-related cysteine endopeptidase 3 Antibody, EC:3.4.22.- Antibody, ATG 4C Antibody, AUT like 1, cysteine endopeptidase Antibody, EC 3.4.22 Antibody, APG4C Antibody, Autophagin-3 Antibody, OTTHUMP00000010715 Antibody, AURL3 Antibody, APG4 autophagy 4 homolog C (*S. cerevisiae*) Antibody, Autophagy related cysteine endopeptidase 3 Antibody, ATG4C Antibody, Autophagin 3 Antibody, ATG4 autophagy related 4 homolog C Antibody, AUT like 3 cysteine endopeptidase Antibody, AURL1 Antibody, ATG4C_HUMAN Antibody, AUT (*S. cerevisiae*) like 1, cysteine endopeptidase Antibody, Autophagy-related protein 4 homolog C Antibody, Autophagy related protein 4 homolog C Antibody, ATG4 autophagy related 4 homolog C (*S. cerevisiae*) Antibody, Autophagy related 4C cysteine peptidase Antibody, Atg4c Antibody, APG4 C Antibody, FLJ14867 Antibody, AUT-like 3 cysteine endopeptidase Antibody,

Immunogen

Synthetic peptide from the C-terminal of Human ATG4C (aa. 296-306)

Purification



Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-ATG4C Polyclonal Antibody (ASM10514). Tissue: Colon carcinoma cell line (RKO). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Rabbit Anti-ATG4C Polyclonal Antibody (ASM10514) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Rabbit ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60 min at RT, 5 min at RT. Localization: Cytoplasm. Magnification: 60X. (A) DAPI nuclear stain. (B) Phalloidin Texas Red F-Actin stain. (C) ATG4C Antibody. (D) Composite.

Peptide Affinity Purified

Storage **-20°C**

Storage Buffer

PBS, 50% glycerol, 0.09% sodium azide

Shipping **Blue Ice or 4°C**

Temperature

Certificate of Analysis

A 1:1000 dilution of SPC-633 was sufficient for detection of ATG4C in 15 µg of Human HeLa Cell Lysates by ECL immunoblot analysis using goat anti-rabbit IgG:HRP as the secondary antibody.

Cellular Localization

Cytoplasm

ATG4C Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

