

RIP3 Antibody
Catalog # **ASM10538**

Specification

RIP3 Antibody - Product Information

Application	ICC/IF, WB
Primary Accession	Q9Y572
Other Accession	NP_006862.2
Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal

Description

Rabbit Anti-Human RIP3 Polyclonal

Target/Specificity

Detects ~57 kDa.

Other Names

RIPK3 Antibody, Receptor-interacting protein 3 Antibody, Receptor interacting protein 3 Antibody, Receptor interacting serine/threonine protein kinase 3 Antibody, RIP-3 Antibody, RIPK3_HUMAN Antibody, Receptor-interacting serine/threonine-protein kinase 3 Antibody, RIPK 3 Antibody, RIP 3 Antibody, Receptor interacting serine threonine kinase 3 Antibody, RIP3 Antibody, RIP like protein kinase 3 Antibody, RIP-like protein kinase 3 Antibody

Immunogen

Synthetic peptide from the N-terminal to middle of Human RIP3

Purification

Peptide Affinity Purified

Storage **-20°C**

Storage Buffer


PBS, 50% glycerol, 0.09% sodium azide

Shipping **Blue Ice or 4°C**
Temperature

Certificate of Analysis

A 1:1000 dilution of SPC-672 was sufficient for detection of RIP3 in 15 µg of mouse brain cell lysates by ECL immunoblot analysis using goat anti-rabbit IgG:HRP as the secondary antibody.

Cellular Localization




Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-RIP3 Polyclonal Antibody (ASM10538). Tissue: Colon carcinoma cell line (RKO). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Rabbit Anti-RIP3 Polyclonal Antibody (ASM10538) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Rabbit ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60 min at RT, 5 min at RT. Localization: Cytoplasm. Magnification: 60X. (A) DAPI nuclear stain. (B) Phalloidin Texas Red F-Actin stain. (C) RIP3 Antibody. (D) Composite.

Cytoplasm | Cytosol | Cell Membrane |
Mitochondrion

RIP3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Western blot analysis of Mouse Brain cell lysates showing detection of ~56.9 kDa RIP3 protein using Rabbit Anti-RIP3 Polyclonal Antibody (ASM10538). Lane 1: Molecular Weight Ladder (MW). Lane 2: Mouse Brain cell lysates. Load: 15 μ g. Block: 5% Skim Milk in 1X TBST. Primary Antibody: Rabbit Anti-RIP3 Polyclonal Antibody (ASM10538) at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Rabbit IgG: HRP at 1:1000 for 60 min at RT. Color Development: ECL solution for 6 min in RT. Predicted/Observed Size: ~56.9 kDa.