

Sheep IgG F(c) Peroxidase

Catalog # ASR1747

Specification

Sheep IgG F(c) Peroxidase - Product Information

Description SHEEP IgG F(c)

fragment Peroxidase conjugated Peroxidase

Conjugate Peroxidase (Horseradish)

Physical State
Host Isotype
Buffer

Lyophilized
IgG F(c)
0.02 M
Potassium

Phosphate, 0.15

M Sodium

Chloride, pH 7.2

Species of Origin
Reconstitution
Sheep
1.0 mL

Volume

Reconstitution

Buffer deionized water

(or equivalent)

Restore with

Stabilizer 10 mg/mL Bovine

Serum Albumin

(BSA) -

Immunoglobulin and Protease

free

Preservative 0.01% (w/v)

Gentamicin Sulfate. Do NOT add Sodium

Azide!

Sheep IgG F(c) Peroxidase - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from normal serum by delipidation, salt fractionation, ion exchange chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Sheep IgG, anti-Sheep IgG F(c) and





Tel: 858.875.1900 Fax: 858.622.0609

anti-Sheep Serum. No reaction was observed against anti-Sheep IgG F(ab')2 or anti-Papain.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Sheep IgG F(c) Peroxidase - Protein Information

Sheep IgG F(c) Peroxidase - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cvtometv
- Cell Culture