

Anti-Armenian Hamster IgG (H&L) (Fluorescein Conjugated) Secondary Antibody **Goat Polyclonal, Fluorescein (FITC)** Catalog # ASR1960

Specification

Anti-Armenian Hamster IgG (H&L) (Fluorescein Conjugated) Secondary Antibody - Product Information

Description Anti-ARMENIAN

HAMSTER IgG (H&L) (GOAT) Antibody **Fluorescein** Conjugated

Host Goat

Fluorescein Conjugate

(FITC)

FP Value 2.8 moles

Fluorescein (FITC)

per mole of IgG

Target Species Armenian

Hamster Hamster **Polyclonal**

,3,4, Application

Reactivity

Clonality

FLISA 1:10,000-1: **Application Note**

50.000:IF

Microscopy 1:1,00

0-1:5,000;FlowCyt

ometry

1:500-1:2,500;

Physical State Lyophilized

Host Isotype laG

Target Isotype IgG (H&L) Buffer 0.02 M

Potassium Phosphate, 0.15

M Sodium

Chloride, pH 7.2 **Armenian Immunogen**

Hamster IgG whole molecule

Restore with

Reconstitution 1.0 mL

Volume


Reconstitution

Buffer

deionized water (or equivalent) Stabilizer 10 mg/mL Bovine Serum Albumin

(BSA) -


Immunoglobulin and Protease



Western Blot of Fluorescein conjugated Goat anti-Armenian Hamster IgG secondary antibody. Lane 1: Armenian Hamster IgG. Lane 2: none. Load: 50 ng per lane. Primary antibody: none. Secondary antibody: Fluorescein goat secondary antibody at 1:1,000 for 60 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 25 & 55 kDa, 25 & 55 kDa for Armenian Hamster IgG. Other band(s): none.

Anti-Armenian Hamster IgG (H&L) (Fluorescein Conjugated) Secondary **Antibody - Background**

FITC Anti-Hamster IgG Secondary Antibody is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.





Preservative

free 0.01% (w/v) Sodium Azide

Anti-Armenian Hamster IgG (H&L) (Fluorescein Conjugated) Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Armenian Hamster IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Goat Serum, Armenian Hamster IgG and Armenian Hamster Serum. Minimal reactivity will occur against Golden Syrian Hamster IgG.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note


This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Armenian Hamster IgG (H&L) (Fluorescein Conjugated) Secondary Antibody - Protein Information

Anti-Armenian Hamster IgG (H&L) (Fluorescein Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence





ImmunoprecipitationFlow CytometyCell Culture