

**F(ab')<sub>2</sub> Anti-Rat IgG F(c) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody**  
 Goat Polyclonal, Rhodamine (TRITC)  
 Catalog # ASR2872

**Specification**

**F(ab')<sub>2</sub> Anti-Rat IgG F(c) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Product Information**

Description	<b>F(ab')<sub>2</sub> Anti-RAT IgG F(c) (GOAT) Antibody Rhodamine Conjugated Min X Bv Hs &amp; Hu Serum Proteins</b>
Host Conjugate	<b>Goat Rhodamine (TRITC)</b>
FP Value	<b>4.0 moles Rhodamine (TRITC) per mole of IgG</b>
Target Species	<b>F(ab')<sub>2</sub> Rat</b>
Clonality	<b>Polyclonal</b>
Application	<b>,3,4,</b>
Application Note	<b>FLISA 1:10,000-1:50,000;IF Microscopy 1:1,000-1:5,000;FlowCytometry 1:500-1:2,500</b>
Physical State	<b>Lyophilized</b>
Host Isotype	<b>IgG F(ab')<sub>2</sub></b>
Target Isotype	<b>IgG F(c)</b>
Buffer	<b>0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>
Immunogen	<b>Rat IgG F(c) fragment</b>
Reconstitution Volume	<b>1.0 mL</b>
Reconstitution Buffer	<b>Restore with deionized water (or equivalent)</b>
Stabilizer	<b>10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease</b>

**F(ab')<sub>2</sub> Anti-Rat IgG F(c) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Background**

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

Preservative                    **free**  
   **0.01% (w/v)**  
   **Sodium Azide**

**F(ab')<sub>2</sub> Anti-Rat IgG F(c) (Rhodamine Conjugated)  
Pre-Adsorbed Secondary Antibody - Additional  
Information**

**Shipping Condition**

Ambient

**Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rat IgG, Rat IgG F(c) and Rat Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Rat IgG F(ab')<sub>2</sub> or Bovine, Horse and Human Serum Proteins.

**Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**F(ab')<sub>2</sub> Anti-Rat IgG F(c) (Rhodamine Conjugated)  
Pre-Adsorbed Secondary Antibody - Protein  
Information**

**F(ab')<sub>2</sub> Anti-Rat IgG F(c) (Rhodamine  
Conjugated) Pre-Adsorbed Secondary  
Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)