

Anti-Mouse IgG F(c) (Peroxidase Conjugated) Secondary Antibody
Goat Polyclonal, Peroxidase (Horseradish)
Catalog # ASR3396

Specification

**Anti-Mouse IgG F(c) (Peroxidase Conjugated)
Secondary Antibody - Product Information**

Description	Anti-MOUSE IgG F(c) (GOAT) Antibody Peroxidase Conjugated Goat Peroxidase (Horseradish)
Host Conjugate	Goat Peroxidase (Horseradish)
Target Species	Mouse
Reactivity	Mouse
Clonality	Polyclonal
Application	,1,2,10,
Application Note	ELISA 1:10,000-1:50,000;Western Blot 1:1,000-1:10,000;Immunohistochemistry 1:500-1:2,500
Physical State	Lyophilized
Host Isotype	IgG
Target Isotype	IgG F(c)
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Anti-Mouse IgG F(c) was produced by repeated immunization with mouse IgG F(c) fragment in goat.
Reconstitution Volume	2.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v)

Anti-Mouse IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Background

Anti-Mouse IgG F(c) antibody generated in goat detects specifically mouse IgG. This secondary antibody anti-Mouse is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays.

**Gentamicin
Sulfate. Do NOT
add Sodium
Azide!**

**Anti-Mouse IgG F(c) (Peroxidase Conjugated)
Secondary Antibody - Additional Information**

Shipping Condition

Ambient

Purity

This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against Mouse IgG F(ab'à2).

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**Anti-Mouse IgG F(c) (Peroxidase Conjugated)
Secondary Antibody - Protein Information**

**Anti-Mouse IgG F(c) (Peroxidase
Conjugated) Secondary Antibody -
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)