

SLC27A4 Antibody (C-term) Blocking Peptide

Synthetic peptide

Catalog # BP13227b

Specification**SLC27A4 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [O6P1M0](#)**SLC27A4 Antibody (C-term) Blocking Peptide - Additional Information**

Gene ID 10999

Other Names

Long-chain fatty acid transport protein 4, FATP-4, Fatty acid transport protein 4, 621-, Solute carrier family 27 member 4, SLC27A4, ACSVL4, FATP4

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13227b was selected from the C-term region of SLC27A4. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

SLC27A4 Antibody (C-term) Blocking Peptide - Protein InformationName SLC27A4 ([HGNC:10998](#))**SLC27A4 Antibody (C-term) Blocking Peptide - Background**

This gene encodes a member of a family of fatty acid transport proteins, which are involved in translocation of long-chain fatty acids across the plasma membrane. This protein is expressed at high levels on the apical side of mature enterocytes in the small intestine, and appears to be the principal fatty acid transporter in enterocytes. Clinical studies suggest this gene as a candidate gene for the insulin resistance syndrome. Mutations in this gene have been associated with ichthyosis prematurity syndrome.

SLC27A4 Antibody (C-term) Blocking Peptide - References

Morice-Picard, F., et al. Am. J. Med. Genet. A 152A (10), 2664-2665 (2010) ; Yokoyama, K., et al. Nephron Clin Pract 115 (4), C237-C243 (2010) ; Ban, H.J., et al. BMC Genet. 11, 26 (2010) ; Klar, J., et al. Am. J. Hum. Genet. 85(2):248-253(2009) ; Jia, Z., et al. J. Mol. Neurosci. 33(1):25-31(2007)

Function

Involved in translocation of long-chain fatty acids (LFCA) across the plasma membrane (PubMed:12556534, PubMed:21395585). Has acyl-CoA ligase activity for long-chain and very-long-chain fatty acids (VLCFAs) (PubMed:24269233). Appears to be the principal fatty acid transporter in small intestinal enterocytes. Plays a role in the formation of the epidermal barrier. Required for fat absorption in early embryogenesis (By similarity). Probably involved in fatty acid transport across the blood barrier (PubMed:21395585). Indirectly inhibits RPE65 via substrate competition and via production of VLCFA derivatives like lignoceroyl-CoA. Prevents light-induced degeneration of rods and cones (By similarity).

Cellular Location

Endoplasmic reticulum membrane;
Multi-pass membrane protein

Tissue Location

Expressed at highest levels in brain, testis, colon and kidney. Expressed at medium levels in heart and liver, small intestine and stomach. Expressed at low levels in peripheral leukocytes, bone marrow, skeletal muscle and aorta. Expressed in adipose tissue (PubMed:24269233, PubMed:9878842). Expressed in brain gray matter (PubMed:21395585).

SLC27A4 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)