

WNT3A Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP14596c**Specification****WNT3A Antibody (Center) Blocking Peptide - Product Information**Primary Accession [P56704](#)**WNT3A Antibody (Center) Blocking Peptide - Additional Information**

Gene ID 89780

Other Names

Protein Wnt-3a, WNT3A

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

WNT3A Antibody (Center) Blocking Peptide - Protein Information

Name WNT3A

Function

Ligand for members of the frizzled family of seven transmembrane receptors (Probable). Functions in the canonical Wnt signaling pathway that results in activation of transcription factors of the TCF/LEF family (PubMed:20093360, PubMed:21244856).

WNT3A Antibody (Center) Blocking Peptide - Background

The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family. It encodes a protein which shows 96% amino acid identity to mouse Wnt3A protein, and 84% to human WNT3 protein, another WNT gene product. This gene is clustered with WNT14 gene, another family member, in chromosome 1q42 region.

WNT3A Antibody (Center) Blocking Peptide - References

Sengupta, P.K., et al. J. Biol. Chem. 285(41):31164-31173(2010) Hubner, R., et al. Biochem. Biophys. Res. Commun. 400(3):358-362(2010) von Marschall, Z., et al. Biochem. Biophys. Res. Commun. 400(3):299-304(2010) Shin, H., et al. J. Dermatol. Sci. 58(2):91-96(2010) Jugessur, A., et al. PLoS ONE 5 (7), E11493 (2010) :

PubMed:24841207, PubMed:26902720). Required for normal embryonic mesoderm development and formation of caudal somites. Required for normal morphogenesis of the developing neural tube (By similarity). Mediates self-renewal of the stem cells at the bottom on intestinal crypts (in vitro) (PubMed:26902720).

Cellular Location

Secreted, extracellular space, extracellular matrix {ECO:0000250|UniProtKB:P27467}. Secreted

Tissue Location

Moderately expressed in placenta and at low levels in adult lung, spleen, and prostate.

WNT3A Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)