

FOLR1 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP18815a**Specification****FOLR1 Antibody (N-term) Blocking Peptide - Product Information**Primary Accession [P15328](#)**FOLR1 Antibody (N-term) Blocking Peptide - Additional Information**

Gene ID 2348

Other Names

Folate receptor alpha, FR-alpha, Adult folate-binding protein, FBP, Folate receptor 1, Folate receptor, adult, KB cells FBP, Ovarian tumor-associated antigen MOv18, FOLR1, FOLR

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

FOLR1 Antibody (N-term) Blocking Peptide - Protein Information

Name FOLR1

Synonyms FOLR

Function

Binds to folate and reduced folic acid derivatives and mediates delivery of 5-methyltetrahydrofolate and folate analogs into the interior of cells. Has high affinity for folate and folic acid analogs at neutral pH.

FOLR1 Antibody (N-term) Blocking Peptide - Background

The protein encoded by this gene is a member of the folatereceptor family. Members of this gene family bind folic acid and its reduced derivatives, and transport 5-methyltetrahydrofolate into cells. This gene product is a secreted protein that either anchors to membranes via a glycosyl-phosphatidylinositol linkage or exists in a soluble form. Mutations in this gene have been associated with neurodegeneration due to cerebral folate transport deficiency. Due to the presence of two promoters, multiple transcription start sites, and alternative splicing, multiple transcript variants encoding the same protein have been found for this gene.

FOLR1 Antibody (N-term) Blocking Peptide - References

Sivakumaran, S., et al. J. Steroid Biochem. Mol. Biol. 122(5):333-340(2010)
Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
O'Byrne, M.R., et al. Birth Defects Res. Part A Clin. Mol. Teratol. 88(8):689-694(2010)
Jugessur, A., et al. PLoS ONE 5 (7), E11493 (2010)
Elwood, P.C., et al. Biochemistry 36(6):1467-1478(1997)

Exposure to slightly acidic pH after receptor endocytosis triggers a conformation change that strongly reduces its affinity for folates and mediates their release. Required for normal embryonic development and normal cell proliferation.

Cellular Location

Cell membrane; Lipid-anchor, GPI-anchor. Secreted. Cytoplasmic vesicle. Cytoplasmic vesicle, clathrin-coated vesicle. Endosome. Apical cell membrane Note=Endocytosed into cytoplasmic vesicles and then recycled to the cell membrane

Tissue Location

Primarily expressed in tissues of epithelial origin. Expression is increased in malignant tissues. Expressed in kidney, lung and cerebellum. Detected in placenta and thymus epithelium.

**FOLR1 Antibody (N-term) Blocking Peptide
- Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)