

GAS2 Antibody (N-term) Blocking Peptide

Synthetic peptide Catalog # BP18918a

Specification

GAS2 Antibody (N-term) Blocking Peptide - Product Information

Primary Accession <u>043903</u>

GAS2 Antibody (N-term) Blocking Peptide - Additional Information

Gene ID 2620

Other Names

Growth arrest-specific protein 2, GAS-2, GAS2

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

GAS2 Antibody (N-term) Blocking Peptide - Protein Information

Name GAS2

Function

May play a role in apoptosis by acting as a cell death substrate for caspases. Is cleaved during apoptosis and the cleaved form induces dramatic rearrangements of the actin cytoskeleton and potent changes in the shape of the affected cells. May be involved in the membrane ruffling process (By similarity).


Cellular Location

GAS2 Antibody (N-term) Blocking Peptide - Background

The protein encoded by this gene is a caspase-3 substratethat plays a role in regulating microfilament and cell shapechanges during apoptosis. It can also modulate cell susceptibilityto p53-dependent apoptosis by inhibiting calpain activity. Multiplealternatively spliced variants, encoding the same protein, havebeen identified.

GAS2 Antibody (N-term) Blocking Peptide - References

Huang, W., et al. Mol. Cell. Biol. 30(19):4575-4594(2010)Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)Benetti, R., et al. J. Biol. Chem. 280(23):22070-22080(2005)Tsutsumi, S., et al. Am. J. Hum. Genet. 74(6):1255-1261(2004)Benetti, R., et al. EMBO J. 20(11):2702-2714(2001)





Cytoplasm, cytoskeleton, stress fiber.
Membrane
{ECO:0000250|UniProtKB:P11862};
Peripheral membrane protein
{ECO:0000250|UniProtKB:P11862}
Note=Component of the microfilament
system. Colocalizes with actin fibers at the
cell border and along the stress fibers in
growth- arrested fibroblasts. Mainly
membrane-associated. When
hyperphosphorylated, accumulates at
membrane ruffles
{ECO:0000250|UniProtKB:P11862}

Tissue Location

Ubiquitously expressed with highest levels in liver, lung, and kidney. Not found in spleen

GAS2 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides