

Rde-1 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP1964c**Specification****Rde-1 Antibody (Center) Blocking Peptide - Product Information**Primary Accession [G5EEH0](#)
Other Accession [Q9XU82](#)**Rde-1 Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 179393**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP1964c](/product/products/AP1964c) was selected from the Center region of human Rde-1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Rde-1 Antibody (Center) Blocking Peptide - Protein Information**Name** rde-1
{ECO:0000313|EMBL:CAB05546.2,
ECO:0000313|WormBase:K08H10.7}**Rde-1 Antibody (Center) Blocking Peptide - Background**

Double-stranded (ds) RNA is a potent sequence-specific inhibitor of gene function. RNA interference (RNAi) is a cellular defense mechanism that uses double-stranded RNA (dsRNA) as a sequence-specific trigger to guide the degradation of homologous single-stranded RNAs. RNAi is a multistep process involving several proteins and at least one type of RNA intermediate, a population of small 21-25 nt RNAs (called siRNAs) that are initially derived from cleavage of the dsRNA trigger. Genetic studies have implicated several RNA interference-deficient (rde) family members in germline maintenance and development, and several simple loss of function mutants have been identified. Family members rde-1 and rde-4 are required for RNAi but are not essential for organismal viability. While rde-1 and rde-4 are distinct from other RNAi-deficient family members both both for their inability to mobilize transposons and lack of chromosome loss, each appears to have a distinct role in the interference mechanism. Evidence indicates that rde-4 is involved before or during production of siRNAs, whereas rde-1 acts after the siRNAs have been formed.

Rde-1 Antibody (Center) Blocking Peptide - References

Tabara H, et al. Cell. 2002. 109(7):861-71.
Parrish S, et al. RNA. 2001. 7(10):1397-402.
Fagard M, et al. PNAS. 2000. 97(21):11650-4
Ketting RF, et al. Nature. 2000.
404(6775):296-8. Tabara H, et al. Cell. 1999.
99(2):123-32.

**Rde-1 Antibody (Center) Blocking Peptide
- Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)