

JUN Antibody (S63) Blocking Peptide
Synthetic peptide
Catalog # BP1984d**Specification****JUN Antibody (S63) Blocking Peptide - Product Information**Primary Accession [P05412](#)**JUN Antibody (S63) Blocking Peptide - Additional Information**

Gene ID 3725

Other Names

Transcription factor AP-1, Activator protein 1, AP1, Proto-oncogene c-Jun, V-jun avian sarcoma virus 17 oncogene homolog, p39, JUN

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP1984d](/product/products/AP1984d) was selected from the S63 region of human JUN. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

JUN Antibody (S63) Blocking Peptide - Protein Information

Name JUN

JUN Antibody (S63) Blocking Peptide - Background

JUN interacts directly with specific target DNA sequences to regulate gene expression. Jun recognizes the AP-1 consensus sequence TGACTCA, a response element that confers sensitivity to one of the tumor-promoting Phorbol esters, 12-O-tetradecanoyl-phorbol-13-acetate (see also: TRE, TPA response element). Jun itself forms homodimers or heterodimers with junD and junB and also interacts with the oncogene product fos, forming jun-fos heterodimers.

JUN Antibody (S63) Blocking Peptide - References

Fujita,S., J. Mol. Biol. 378 (3), 492-504 (2008)
Gan,X.Q., . Cell Biol. 180 (6), 1087-1100 (2008)
Yogev,O., Cancer Res. 68 (5), 1398-1406 (2008)

Function

Transcription factor that recognizes and binds to the enhancer heptamer motif 5'-TGA[CG]TCA-3' (PubMed:10995748, PubMed:22083952). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation (PubMed:17210646). Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells (PubMed:24623306). Binds to the USP28 promoter in colorectal cancer (CRC) cells (PubMed:24623306).

Cellular Location

Nucleus.

Tissue Location

Expressed in the developing and adult prostate and prostate cancer cells.

JUN Antibody (S63) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)