



TSK Antibody (Center) Blocking peptide

Synthetic peptide Catalog # BP5987c

Specification

TSK Antibody (Center) Blocking peptide - Product Information

Primary Accession <u>Q8WUA8</u> Other Accession <u>NP 056331.2</u>

TSK Antibody (Center) Blocking peptide - Additional Information

Gene ID 25987

Other Names

Tsukushin, Tsukushi, E2-induced gene 4 protein, Leucine-rich repeat-containing protein 54, TSKU, E2IG4, LRRC54, TSK

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

TSK Antibody (Center) Blocking peptide - Protein Information

Name TSKU

Synonyms E2IG4, LRRC54, TSK

Function

Contributes to various developmental events and other processes such as wound healing and cholesterol homeostasis through its interactions with multiple signaling pathways. Wnt signaling inhibitor which competes with WNT2B for binding to



Wnt receptor FZD4 and represses WNT2B-dependent development of the peripheral eye. Plays a role in regulating the hair cycle by controlling TGFB1 signaling. Required for the development of the anterior commissure in the brain by inhibiting neurite outgrowth. Essential for terminal differentiation of hippocampal neural stem cells. Plays a role in regulating bone elongation and bone mass by modulating growth plate chondrocyte function and overall body size. Required for development of the inner ear through its involvement in stereocilia formation in inner hair cells. Facilitates wound healing by inhibiting secretion of TGFB1 from macrophages which prevents myofibroblast differentiation, maintaining inflammatory cell quiescence. Plays a role in cholesterol homeostasis by reducing circulating high-density lipoprotein cholesterol, lowering cholesterol efflux capacity and decreasing cholesterol-to-bile acid conversion in the liver. In one study, shown to negatively regulate sympathetic innervation in brown fat, leading to reduced energy expenditure. In another study, shown not to affect brown fat thermogenic capacity, body weight gain or glucose homeostasis.

Cellular LocationSecreted
{ECO:0000250|UniProtKB:Q8CBR6}.

TSK Antibody (Center) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides