

MLK4 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP8006c**Specification****MLK4 Antibody (Center) Blocking Peptide -
Product Information**Primary Accession [Q5TCX8](#)**MLK4 Antibody (Center) Blocking Peptide -
Additional Information**

Gene ID 84451

Other NamesMitogen-activated protein kinase kinase
kinase MLK4, Mixed lineage kinase 4, MLK4
{ECO:0000303|Ref1}**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP8006c](/product/products/AP8006c) was selected from the Center region of human MLK4. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**MLK4 Antibody (Center) Blocking Peptide -
Protein Information**Name MAP3K21 ([HGNC:29798](#))**MLK4 Antibody (Center) Blocking Peptide -
Background**

Mixed-lineage kinases (MLKs) function as mitogen-activated protein kinase kinase kinases (MAP3Ks), activating the p38 and JNK signaling cascades. Systematic mutational analyses of tyrosine kinases reveal that a minimum of 30% of colorectal cancer contain at least one mutation in the tyrosine kinases; sampling of colorectal cancer cells suggest an important role for MLK4 in this process.

**MLK4 Antibody (Center) Blocking Peptide -
References**

Gregory,S.G., Nature 441 (7091), 315-321
(2006)

Function

Negative regulator of TLR4 signaling. Does not activate JNK1/MAPK8 pathway, p38/MAPK14, nor ERK2/MAPK1 pathways.

MLK4 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)