

ULK3 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP8115c**Specification**

**ULK3 Antibody (Center) Blocking Peptide -
Product Information**Primary Accession [Q6PHR2](#)**ULK3 Antibody (Center) Blocking Peptide -
Additional Information**

Gene ID 25989

Other NamesSerine/threonine-protein kinase ULK3,
Unc-51-like kinase 3, ULK3**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP8115c](/product/products/AP8115c) was selected from the Center region of human ULK3. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**ULK3 Antibody (Center) Blocking Peptide -
Protein Information**

Name ULK3

Function**ULK3 Antibody (Center) Blocking Peptide -
Background**

ULK3 is a member of the ULK kinase family, one of three human homologs of the yeast autophagy-specific kinase.

**ULK3 Antibody (Center) Blocking Peptide -
References**

The MGC Project Team, Genome Res. 14:2121-2127(2004).

Serine/threonine protein kinase that acts as a regulator of Sonic hedgehog (SHH) signaling and autophagy. Acts as a negative regulator of SHH signaling in the absence of SHH ligand: interacts with SUFU, thereby inactivating the protein kinase activity and preventing phosphorylation of GLI proteins (GLI1, GLI2 and/or GLI3). Positively regulates SHH signaling in the presence of SHH: dissociates from SUFU, autophosphorylates and mediates phosphorylation of GLI2, activating it and promoting its nuclear translocation. Phosphorylates in vitro GLI2, as well as GLI1 and GLI3, although less efficiently. Also acts as a regulator of autophagy: following cellular senescence, able to induce autophagy.

Cellular Location

Cytoplasm. Note=Localizes to pre-autophagosomal structure during cellular senescence

Tissue Location

Widely expressed. Highest levels observed in fetal brain. In adult tissues, high levels in brain, liver and kidney, moderate levels in testis and adrenal gland and low levels in heart, lung, stomach, thymus, prostate and placenta. In the brain, highest expression in the hippocampus, high levels also detected in the cerebellum, olfactory bulb and optic nerve. In the central nervous system, lowest levels in the spinal cord

ULK3 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)